

B9



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C07D 259/00, C07F 9/6524, 13/00, 15/02		A1	(11) International Publication Number: WO 96/40658
			(43) International Publication Date: 19 December 1996 (19.12.96)
(21) International Application Number: PCT/US96/07553		(74) Agents: WILLIAMS, Roger, A. et al.; G.D. Searle & Co., Corporate Patent Dept., P.O. Box 5110, Chicago, IL 60680- 5110 (US).	
(22) International Filing Date: 30 May 1996 (30.05.96)			
(30) Priority Data: 08/486,434 7 June 1995 (07.06.95) US		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(60) Parent Application or Grant (63) Related by Continuation US 08/486,434 (CON) Filed on 7 June 1995 (07.06.95)			
(71) Applicant (for all designated States except US): MONSANTO COMPANY [US/US]; 800 North Lindbergh Boulevard, St. Louis, MO 63167 (US).		Published With international search report.	
(72) Inventors; and (75) Inventors/Applicants (for US only): LENNON, Patrick, J. [US/US]; 7540 Wydown Boulevard #3W, Clayton, MI 63105 (US). HENKE, Susan, L. [US/US]; 123 Parsons Avenue, Webster, MO 63119 (US). ASTON, Karl, W. [US/US]; 19040 Sunflower Ridge Lane, Pacific, MO 63069 (US).			
(54) Title: PROCESS FOR PREPARING SUBSTITUTED POLYAZAMACROCYCLES			
(57) Abstract A process for preparing a substituted polyazamacrocycle is provided which comprises contacting a diamine or triamine and a dicarboxylic acid or ester or anhydride thereof in the presence of a suitable base and a suitable solvent to produce the substituted polyazamacrocycle provided that when an ester of said dicarboxylic acid is used, said suitable base is optional, and when said dicarboxylic acid or an anhydride of said dicarboxylic acid is used, the reaction mixture further comprises a suitable coupling agent.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam


PROCESS FOR PREPARING SUBSTITUTED POLYAZAMACROCYCLES

BACKGROUND OF THE INVENTION

- This invention relates to a process for preparing substituted polyazamacrocycles. In one aspect, this invention relates to a process for preparing substituted fifteen-membered and sixteen-membered pentaazamacrocyclic ligands which are useful in the sequestration of transition metal ions. In a further aspect, this invention relates to a process for preparing manganese(II) or manganese(III) complexes of substituted fifteen-membered and sixteen-membered pentaazamacrocyclic ligands which catalytically dismutate superoxide.

- The substituted polyazamacrocycles of the invention, which are useful in the sequestration of transition metal ions, can be used in qualitative or quantitative assays of transition metal ions. The substituted polyazamacrocycles of the invention are also useful as intermediates in the preparation of manganese complexes of substituted fifteen-membered and sixteen-membered pentaazamacrocyclic ligands. The manganese complexes of substituted fifteen-membered and sixteen-membered pentaazamacrocyclic ligands are low molecular weight mimics of superoxide dismutase (SOD) which are useful as therapeutic agents for inflammatory disease states or disorders which are mediated, at least in part, by superoxide. The manganese (II) complexes of substituted fifteen-membered pentaazamacrocyclic ligands are also useful as magnetic resonance imaging (MRI) contrast agents.

- The enzyme superoxide dismutase catalyzes the conversion of superoxide into oxygen and hydrogen peroxide according to equation (1) (hereinafter referred to as dismutation). Reactive oxygen metabolites derived from superoxide are postulated to contribute to the tissue pathology in a number of



- inflammatory diseases and disorders, such as reperfusion injury to the ischemic myocardium, inflammatory bowel disease, rheumatoid arthritis, osteoarthritis, atherosclerosis, hypertension, metastasis, psoriasis, organ transplant rejections, radiation-induced injury, asthma, influenza, stroke, burns and trauma. See, for example, Bulkley, G.B., Reactive oxygen metabolites and reperfusion injury: aberrant triggering of reticuloendothelial function, *The Lancet*, Vol. 344, pp. 934-36, October 1, 1994; Grisham, M.B., Oxidants and free radicals in inflammatory bowel disease, *The Lancet*, Vol. 344, pp. 859-861, September 24, 1994; Cross, C.E. et al., Reactive oxygen species and the lung, *The Lancet*, Vol. 344, pp. 930-33, October 1, 1994; Jenner, P., Oxidative damage in neurodegenerative disease, *The Lancet*, Vol.

-2-

344, pp. 796-798, September 17, 1994; Cerutti, P.A., Oxy-radicals and cancer, *The Lancet*, Vol. 344, pp. 862-863, September 24, 1994 Simic, M. G., et al, Oxygen Radicals in Biology and Medicine, Basic Life Sciences, Vol. 49, Plenum Press, New York and London, 1988; Weiss J. Cell. Biochem., 1991 Suppl. 15C, 216 Abstract C110 (1991); Petkau, A., Cancer Treat. Rev. **13**, 17 (1986); McCord, J. Free Radicals Biol. Med., **2**, 307 (1986); and Bannister, J.V. et al, Crit. Rev. Biochem., **22**, 111 (1987). The above-identified references from *The Lancet* teach the nexus between free radicals derived from superoxide and a variety of diseases. In particular, the Bulkley and Grisham references specifically teach that there is a nexus between the dismutation of superoxide and the final disease treatment.

It is also known that superoxide is involved in the breakdown of endothelium-derived vascular relaxing factor (EDRF), which has been identified as nitric oxide (NO), and that EDRF is protected from breakdown by superoxide dismutase. This suggests a central role for activated oxygen species derived from superoxide in the pathogenesis of vasospasm, thrombosis and atherosclerosis. See, for example, Gryglewski, R.J. et al., "Superoxide Anion is Involved in the Breakdown of Endothelium-derived Vascular Relaxing Factor", *Nature*, Vol. 320, pp. 454-56 (1986) and Palmer, R.M.J. et al., "Nitric Oxide Release Accounts for the Biological Activity of Endothelium Derived Relaxing Factor", *Nature*, Vol. 327, pp. 523-26 (1987).

Clinical trials and animal studies with natural, recombinant and modified superoxide dismutase enzymes have been completed or are ongoing to demonstrate the therapeutic efficacy of reducing superoxide levels in the disease states noted above. However, numerous problems have arisen with the use of the enzymes as potential therapeutic agents, including lack of oral activity, short half-lives *in vivo*, immunogenicity with nonhuman derived enzymes, and poor tissue distribution.

SUMMARY OF THE INVENTION

It is an object of the invention to provide a process for preparing substituted fifteen-membered and sixteen-membered pentaazamacrocyclic ligands. It is a further object of the invention to provide a process for preparing reduced substituted fifteen-membered and sixteen-membered pentaazamacrocyclic ligands. It is yet a further object of the invention to provide a process for preparing manganese(II) or manganese(III) complexes of substituted fifteen-membered and sixteen-membered pentaazamacrocyclic ligands.

According to the invention, a process for preparing a substituted polyazamacrocyclic is provided which comprises contacting a diamine and a dicarboxylic acid or ester or anhydride thereof in the presence of a suitable base and a


-3-

suitable solvent under reaction conditions sufficient to produce the substituted polyazamacrocycle, provided that when an ester of said dicarboxylic acid is used, said suitable base is optional and when said dicarboxylic acid or an anhydride of said dicarboxylic acid is used, the reaction mixture further comprises a suitable coupling agent. In one embodiment of the invention, the substituted polyazamacrocycle prepared according to the invention is reduced to produce a reduced substituted polyazamacrocycle. In a further embodiment of the invention, a manganese complex of the reduced substituted polyazamacrocycle is produced by reacting the reduced substituted polyazamacrocycle with a manganese compound under essentially anhydrous and anaerobic conditions and, optionally, further conducting an exchange reaction with the manganese complex to exchange the axial ligands on the manganese complex.

Further according to the invention, a process for preparing a substituted polyazamacrocycle is provided which comprises contacting a triamine and a dicarboxylic acid or ester or anhydride thereof in the presence of a suitable base and a suitable solvent under conditions of time and temperature sufficient to produce the substituted polyazamacrocycle, provided that when an ester of said dicarboxylic acid is used, said suitable base is optional and when said dicarboxylic acid or an anhydride of said dicarboxylic acid is used, the reaction mixture further comprises a suitable coupling agent. In one embodiment of the invention, the substituted polyazamacrocycle prepared according to the invention is reduced to produce a reduced substituted polyazamacrocycle. In a further embodiment of the invention, a manganese complex of the reduced substituted polyazamacrocycle is produced by reacting the reduced substituted polyazamacrocycle with a manganese compound under essentially anhydrous and anaerobic conditions and, optionally, further conducting an exchange reaction with the manganese complex to exchange the axial ligands on the manganese complex.


DETAILED DESCRIPTION OF THE INVENTION

One aspect of this invention relates to a process for preparing a substituted polyazamacrocycle comprising contacting (a) a diamine represented by the formula:




-4-

and (b) a dicarboxylic acid or ester thereof represented by the formula:



- 5 or an anhydride thereof; in the presence of a suitable base and a suitable solvent under reaction conditions sufficient to produce the substituted polyazamacrocyclic represented by the formula:



- provided that when an ester of said dicarboxylic acid is used, said suitable base is optional and when said dicarboxylic acid or an anhydride of said dicarboxylic acid is used, the reaction mixture further comprises a suitable coupling agent; wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₁₁ and R₁₂ are independently selected from hydrogen or alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, alkylcycloalkyl, alkenylcycloalkyl,

-5-

- alkylcycloalkenyl, alkenylcycloalkenyl, heterocyclic, aryl and aralkyl radicals or radicals attached to the α -carbon of α -amino acids; or
- R_3 or R_4 and R_5 or R_6 , and R_7 or R_8 and R_9 or R_{10} together with the carbon atoms to which they are attached independently form a saturated, partially saturated or
- 5 unsaturated cyclic having 3 to 20 carbon atoms; or
- R_1 or R_2 and R_3 or R_4 , R_5 or R_6 and R_7 or R_8 , and R_9 or R_{10} and R_{11} or R_{12} together with the carbon atoms to which they are attached independently form a nitrogen containing heterocycle having 2 to 20 carbon atoms provided that when the nitrogen containing heterocycle is an aromatic heterocycle which does not contain a
- 10 substituent attached to the nitrogen, the hydrogen attached to the nitrogen in the formula, which nitrogen is also in the macrocycle and the R groups attached to the same carbon atoms of the macrocycle are absent;
- R_1 and R_2 , R_3 and R_4 , R_5 and R_6 , R_7 and R_8 , R_9 and R_{10} , and R_{11} and R_{12} together with the carbon atom to which they are attached independently form a saturated,
- 15 partially saturated or unsaturated ring structure having 3 to 20 carbon atoms, or R_1 and R_2 , R_3 and R_4 , R_5 and R_6 , R_7 and R_8 , R_9 and R_{10} , and R_{11} and R_{12} independently are =O or =S; and combinations thereof; and
- wherein R_{13} , R_{14} , R_{15} , R_{16} , R_{19} and R_{20} are independently selected from hydrogen or alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cycloalkylalkyl,
- 20 cycloalkylcycloalkyl, cycloalkenylalkyl, alkylcycloalkyl, alkenylcycloalkyl, alkylcycloalkenyl, alkenylcycloalkenyl, heterocyclic, aryl and aralkyl radicals; or
- R_{13} and R_{14} , R_{15} and R_{16} , and R_{19} and R_{20} together with the carbon atom to which they are attached independently form a saturated, partially saturated or unsaturated ring structure having 3 to 20 carbon atoms; or
- 25 R_{19} and R_{20} are independently -OR₂₃, -OH, -SR₂₃, -NR₂₃R₂₄, -P(O)(OR₂₅)(OR₂₆) or -P(O)(R₂₅)(OR₂₆); or
- R_{19} and R_{20} are =O, =S, =NR₂₃, =N-OH, =N-OR₂₃, =N-O-C(O)-R₂₃, or =CR₂₃R₂₄; or
- R_{13} or R_{14} and R_{15} or R_{16} together with the carbon atoms to which they are attached
- 30 independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms, provided that if said diamine has three carbons between the nitrogen atoms, the saturated, partially saturated or unsaturated cyclic has 4 to 20 carbon atoms; or

-6-

R₁₃ or R₁₄ and R₁₉ or R₂₀, or R₁₅ or R₁₆ and R₁₉ or R₂₀ together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms; and combinations thereof; wherein R₁₇ and R₁₈ are independently selected from hydrogen and alkyl or aryl

5 groups;

wherein R is hydrogen or alkyl or aryl groups;


wherein R₂₃ and R₂₄ are independently selected from alkyl, aralkyl or aryl groups;

R₂₅ and R₂₆ are independently selected from hydrogen, alkyl, aralkyl or aryl groups and

- 10 wherein A, B and C are independently selected from hydrogen, alkyl, aryl, aralkyl, cycloalkyl, -OR₂₁, -SO₂R₂₁, -COOR₂₁, -COR₂₁, -CONR₂₁R₂₂, R₂₁R₂₂P(O), (R₂₁O)(R₂₂O)P(O), R₂₁R₂₂P(S), -SOR₂₁ or -Si(OR₂₁)₃, provided that when the two "R" groups on a carbon adjacent to the nitrogen are =O or =S, A, B and C are hydrogen, alkyl, aralkyl or aryl and R₂₁ and R₂₂ are independently selected from
- 15 hydrogen, alkyl, aryl, aralkyl or alkaryl groups.

As used herein, the term "anhydride" means an intramolecular anhydride of the dicarboxylic acid or intermolecular anhydrides of the dicarboxylic acid of linear or cyclic form.

- 20 For producing a reduced substituted polyazamacrocyclic, the process of the invention further comprises reducing the substituted polyazamacrocyclic prepared according to the process of the invention to produce a reduced substituted polyazamacrocyclic of the formula:





-7-

wherein A, B and C are independently selected from hydrogen, alkyl, aryl, aralkyl and cycloalkyl groups, and any "R" groups attached to the same carbon atom of the macrocycle which were =O or =S will be reduced to -OH or -SH.


- 5 The reduction of the substituted polyazamacrocyclic is conducted in the presence of any suitable reducing agent under conventional reduction conditions known to those skilled in the art. The currently preferred reducing agents are selected from the group consisting of aluminum hydrides and boron hydrides. In particular, the preferred reducing agents are selected from the group consisting of lithium
- 10 aluminum hydride, borane and sodium bis(2-methoxyethoxy) aluminum hydride.

- For producing a manganese complex of the reduced substituted polyazamacrocyclic, the process of the invention further comprises reacting the reduced substituted polyazamacrocyclic with a manganese compound under essentially anhydrous and anaerobic conditions to produce a manganese complex represented by
- 15 the formula:



and, optionally, further conducting an exchange reaction with the manganese complex
5 to exchange the ligands X, Y and Z on the manganese complex.


When the diamine is represented by the formula:



-9-


- it is currently preferred that R_{13} , R_{14} , R_{15} and R_{16} are independently selected from hydrogen or alkyl; or R_{13} or R_{14} and R_{15} or R_{16} together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms. In one preferred embodiment, at least one of R_{13} , R_{14} , R_{15} and R_{16} is an alkyl group. More preferably, at least one of R_{13} and R_{14} and at least one of R_{15} and R_{16} is an alkyl group. In another preferred embodiment, R_{13} and R_{16} together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms and R_{14} and R_{15} are hydrogen.

- 10 When the diamine is represented by the formula:




- it is currently preferred that R_{13} , R_{14} , R_{15} , R_{16} , R_{19} and R_{20} are independently selected from hydrogen or alkyl; or R_{13} or R_{14} and R_{19} or R_{20} , or R_{15} or R_{16} and R_{19} or R_{20} together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms; or R_{19} and R_{20} together are $=O$ or $=S$. In a preferred embodiment, at least one of R_{13} or R_{14} and R_{19} or R_{20} , or R_{15} or R_{16} and R_{19} or R_{20} together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms and the remaining "R" groups of the diamine are independently selected from hydrogen or alkyl, preferably hydrogen.

- 20 The currently preferred "R" groups of the dicarboxylic acids or esters thereof of the formula:




-10-

- or anhydrides thereof; are as follows. It is currently preferred that $R_1, R_2, R_3, R_4, R_5, R_6, R_7, R_8, R_9, R_{10}, R_{11}$ and R_{12} are independently selected from hydrogen, alkyl radicals or radicals attached to the α -carbon of α -amino acids; or R_3 or R_4 and R_5 or R_6 , and R_7 or R_8 and R_9 or R_{10} together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms; or R_1 and R_2, R_3 and R_4, R_5 and R_6, R_7 and R_8, R_9 and R_{10} , and R_{11} and R_{12} independently are $=O$ or $=S$. When R_1 and R_2, R_3 and R_4, R_5 and R_6, R_7 and R_8, R_9 and R_{10} , and R_{11} and R_{12} independently are $=O$ or $=S$, it is preferred that each nitrogen atom have no more than one adjacent carbon atom which is substituted with $=O$ or $=S$. In one preferred embodiment, at least one of R_3 or R_4 and R_5 or R_6 , and R_7 or R_8 and R_9 or R_{10} together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms, and the remaining of $R_1, R_2, R_3, R_4, R_5, R_6, R_7, R_8, R_9, R_{10}, R_{11}$ and R_{12} are independently selected from hydrogen, alkyl radicals or radicals attached to the α -carbon of α -amino acids. In another preferred embodiment, at least one, and more preferably at least two, of $R_1, R_2, R_3, R_4, R_5, R_6, R_7, R_8, R_9, R_{10}, R_{11}$ and R_{12} are alkyl radicals or radicals attached to the α -carbon of α -amino acids, and the remaining of $R_1, R_2, R_3, R_4, R_5, R_6, R_7, R_8, R_9, R_{10}, R_{11}$ and R_{12} are hydrogen.
- In either diamine, it is currently preferred that R_{17} and R_{18} are hydrogen. In the dicarboxylic acid, it is currently preferred that R is hydrogen.
- Another aspect of this invention relates to a process for preparing a substituted polyazamacrocyclic comprising contacting (a) a triamine represented by the formula:




and (b) a dicarboxylic acid or ester thereof represented by the formula:

-11-



or an anhydride thereof; in the presence of a suitable base and a suitable solvent under reaction conditions sufficient to produce the substituted polyazamacrocycle represented by the formula:



5

or

- provided that when an ester of the dicarboxylic acid is used, the suitable base is optional and when the dicarboxylic acid or an anhydride of the dicarboxylic acid is used, the reaction mixture further comprises a suitable coupling agent; wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 and R_8 are independently selected from hydrogen or alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, alkylcycloalkyl, alkenylcycloalkyl, alkylcycloalkenyl, alkenylcycloalkenyl, heterocyclic, aryl and aralkyl radicals or radicals attached to the α -carbon of α -amino acids; or
- R_3 or R_4 and R_5 or R_6 together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms; or

-12-

- R_1 or R_2 and R_3 or R_4 , and R_5 or R_6 and R_7 or R_8 together with the carbon atoms to which they are attached independently form a nitrogen containing heterocycle having 2 to 20 carbon atoms provided that when the nitrogen containing heterocycle is an aromatic heterocycle which does not contain a substituent attached to the nitrogen, the hydrogen attached to the nitrogen in the formula, which nitrogen is also in the macrocycle and the R groups attached to the same carbon atoms of the macrocycle are absent; and combinations thereof;
- R_1 and R_2 , R_3 and R_4 , R_5 and R_6 , and R_7 and R_8 together with the carbon atom to which they are attached independently form a saturated, partially saturated or unsaturated ring structure having 3 to 20 carbon atoms or R_1 and R_2 , R_3 and R_4 , R_5 and R_6 , and R_7 and R_8 independently are =O or =S; and combinations thereof; and wherein R_9 , R_{10} , R_{11} , R_{12} , R_{13} , R_{14} , R_{15} , R_{16} , R_{19} and R_{20} are independently selected from hydrogen or alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, alkylcycloalkyl, alkenylcycloalkyl, alkylcycloalkenyl, alkenylcycloalkenyl, heterocyclic, aryl and aralkyl radicals; or
- R_9 or R_{10} and R_{11} or R_{12} , and R_{13} or R_{14} and R_{15} or R_{16} together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms, provided that if R_9 or R_{10} and R_{11} or R_{12} together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic and the diamine has three carbons between the nitrogen atoms, the saturated, partially saturated or unsaturated cyclic has 4 to 20 carbon atoms; or
- R_9 or R_{10} and R_{19} or R_{20} , or R_{11} or R_{12} and R_{19} or R_{20} together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms; or
- R_{11} or R_{12} and R_{13} or R_{14} together with the carbon atoms to which they are attached independently form a nitrogen containing heterocycle having 2 to 20 carbon atoms provided that when the nitrogen containing heterocycle is an aromatic heterocycle which does not contain a substituent attached to the nitrogen, the hydrogen attached to the nitrogen in the formula, which nitrogen is also in the macrocycle and the R groups attached to the same carbon atoms of the macrocycle are absent;
- R_9 and R_{10} , R_{11} and R_{12} , R_{13} and R_{14} , R_{15} and R_{16} , and R_{19} and R_{20} together with the carbon atom to which they are attached independently form a saturated, partially saturated or unsaturated ring structure having 3 to 20 carbon atoms; or

-13-

R_{11} and R_{12} , and R_{13} and R_{14} independently are $=O$ or $=S$; R_{19} and R_{20} independently are $=O$, $=S$, $=NR_{23}$, $=N-OH$, $=N-OR_{23}$, $=N-O-C(O)-R_{23}$, or $=CR_{23}R_{24}$; and combinations thereof;

wherein R_{17} and R_{18} are independently selected from hydrogen and alkyl groups;


5 wherein R is hydrogen or an alkyl group; and

wherein A , B and C are independently selected from hydrogen, alkyl, aryl, aralkyl, cycloalkyl, $-OR_{21}$, $-SO_2R_{21}$, $-COOR_{21}$, $-COR_{21}$, $-CONR_{21}R_{22}$, $R_{21}R_{22}P(O)$, $(R_{21}O)(R_{22}O)P(O)$, $R_{21}R_{22}P(S)$, $-SOR_{21}$ or $-Si(OR_{21})_3$, provided that when the two "R" groups on a carbon adjacent to the nitrogen are $=O$ or $=S$, A , B and C are

10 hydrogen, alkyl, aralkyl or aryl, and R_{21} and R_{22} are independently selected from hydrogen, alkyl, aryl, aralkyl or alkaryl groups.

For producing a reduced substituted polyazamacrocyclic, the process of the invention further comprises reducing the substituted polyazamacrocyclic prepared according to the process of the invention to produce a reduced substituted

15 polyazamacrocyclic of the formula:




wherein A , B and C are independently selected from hydrogen, alkyl, aryl, aralkyl and cycloalkyl groups.

20

For producing a manganese complex of the reduced substituted polyazamacrocyclic, the process of the invention further comprises reacting the reduced substituted polyazamacrocyclic with a manganese compound under essentially


-14-

anhydrous and anaerobic conditions to produce a manganese complex represented by the formula:



5

or




and, optionally, further conducting an exchange reaction with the manganese complex to exchange the ligands X, Y and Z on the manganese complex.

10


-15-

When the triamine is represented by the formula:



- it is currently preferred that R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅ and R₁₆ are independently selected from hydrogen or alkyl; or R₉ or R₁₀ and R₁₁ or R₁₂, and R₁₃ or R₁₄ and R₁₅ or R₁₆ together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms; or R₁₁ and R₁₂, and R₁₃ and R₁₄ independently are =O or =S. It is preferred that only one of R₁₁ and R₁₂, and R₁₃ and R₁₄ are =O or =S. In one preferred embodiment, at least one of R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅ and R₁₆ is an alkyl group. More preferably, at least one of R₁₃ and R₁₄ and at least one of R₁₅ and R₁₆ is an alkyl group or at least one of R₉ and R₁₀ and at least one of R₁₁ and R₁₂ is an alkyl group. In another preferred embodiment, R₉ and R₁₂ or R₁₃ and R₁₆ together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms and the remaining R groups of the triamine are hydrogen.

When the triamine is represented by the formula:




- it is currently preferred that R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₉ and R₂₀ are independently selected from hydrogen or alkyl; or R₉ or R₁₀ and R₁₉ or R₂₀, R₁₁ or R₁₂ and R₁₉ or R₂₀ or R₁₃ or R₁₄ and R₁₅ or R₁₆ together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms; or R₁₁ and R₁₂, R₁₃ and R₁₄, and R₁₉ and R₂₀ independently are =O or =S. It is preferred that only one of R₁₁ and

-16-

- R_{12} , and R_{13} and R_{14} are =O or =S. In a preferred embodiment, at least one of R_9 or R_{10} and R_{19} or R_{20} , R_{11} or R_{12} and R_{19} or R_{20} or R_{13} or R_{14} and R_{15} or R_{16} together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms and the remaining "R" groups of the triamine are independently selected from hydrogen or alkyl, preferably hydrogen.

The currently preferred "R" groups of the dicarboxylic acids or esters thereof of the formula:



- or anhydrides thereof are as follows. It is currently preferred that R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 and R_8 are independently selected from hydrogen, alkyl radicals or radicals attached to the α -carbon of α -amino acids; or R_3 or R_4 and R_5 or R_6 together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms; or R_1 and R_2 , R_3 and R_4 , R_5 and R_6 , and R_7 and R_8 independently are =O or =S. When R_1 and R_2 , R_3 and R_4 , R_5 and R_6 , and R_7 and R_8 independently are =O or =S, it is preferred that each nitrogen atom have no more than one adjacent carbon atom which is substituted with =O or =S. In one preferred embodiment, R_3 or R_4 and R_5 or R_6 together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms, and the remaining of R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 and R_8 are independently selected from hydrogen, alkyl radicals or radicals attached to the α -carbon of α -amino acids. In another preferred embodiment, at least one, and more preferably at least two, of R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 and R_8 are alkyl radicals or radicals attached to the α -carbon of α -amino acids, and the remaining of R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 and R_8 are hydrogen.


In either triamine, it is currently preferred that R_{17} and R_{18} are hydrogen. In the dicarboxylic acid, it is currently preferred that R is hydrogen.

- X, Y and Z represent suitable ligands or charge-neutralizing anions which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof (for example benzoic acid or benzoate anion, phenol or phenoxide anion, alcohol or alkoxide anion). X, Y and Z are independently selected from the group consisting of halide, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxy, hydroperoxy, alkylperoxy, arylperoxy, ammonia, alkylamino, arylamino, heterocycloalkyl amino, heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine, nitric oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol thiocarboxylic acid, alkyl carboxylic acid (such as acetic acid, trifluoroacetic acid, oxalic acid), aryl carboxylic acid (such as benzoic acid, phthalic acid), urea, alkyl urea, aryl urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea, sulfate, sulfite, bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate, thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkyl aryl thiocarbamate, alkyl dithiocarbamate, aryl dithiocarbamate, alkyl aryl dithiocarbamate, bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite, hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluorophosphate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate, tetraaryl borate, tetraalkyl borate, tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino acid, hydroxamic acid, thiotosylate, and anions of ion exchange resins, or systems where one or more of X, Y and Z are independently attached to one or more of the "R" groups, wherein n is an integer from 0 to 3. The preferred ligands from which X, Y and Z are selected include halide, organic acid, nitrate and bicarbonate anions.


-18-

As used herein, "R" groups means all of the R groups attached to the carbon atoms of the polyazamacrocyclic, i.e., R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₉ and R₂₀.


- As utilized herein, the term "alkyl", alone or in combination, means a
 5 straight-chain or branched-chain alkyl radical containing from 1 to about 22 carbon atoms, preferably from about 1 to about 18 carbon atoms, and most preferably from about 1 to about 12 carbon atoms which optionally carries one or more substituents selected from (1) -NR₃₀R₃₁ wherein R₃₀ and R₃₁ are independently selected from hydrogen, alkyl, aryl or aralkyl; or R₃₀ is hydrogen, alkyl, aryl or aralkyl and R₃₁ is
 10 selected from the group consisting of -NR₃₂R₃₃, -OH, -OR₃₄,



- wherein R₃₂ and R₃₃ are independently hydrogen, alkyl, aryl or acyl, R₃₄ is alkyl, aryl or alkaryl, Z' is hydrogen, alkyl, aryl, alkaryl, -OR₃₄, -SR₃₄ or -NR₄₀R₄₁ wherein R₄₀ and R₄₁ are independently selected from hydrogen, alkyl, aryl or
 15 alkaryl, Z'' is alkyl, aryl, alkaryl, -OR₃₄, -SR₃₄ or -NR₄₀R₄₁, R₃₅ is alkyl, aryl, -OR₃₄, or -NR₄₀R₄₁, R₃₆ is alkyl, aryl or -NR₄₀R₄₁, R₃₇ is alkyl, aryl or alkaryl, X' is oxygen or sulfur, and R₃₈ and R₃₉ are independently selected from hydrogen, alkyl or aryl;
 (2) -SR₄₂ wherein R₄₂ is hydrogen, alkyl, aryl, alkaryl, -SR₃₄, -NR₃₂R₃₃,





wherein R₄₃ is -OH, -OR₃₄ or -NR₃₂R₃₃, and A and B are independently -OR₃₄, -SR₃₄ or -NR₃₂R₃₃;




- wherein x is 1 or 2, and R₄₄ is alkyl, aryl, alkaryl, -OH, -OR₃₄, -SR₃₄ or -NR₃₂R₃₃;
 25 (4) -OR₄₅ wherein R₄₅ is hydrogen, alkyl, aryl, alkaryl, -NR₃₂R₃₃,

-19-



wherein R₄₆ is -OH, -SH, -OR₃₄, -SR₃₄ or -NR₃₂R₃₃; or

- 5 (6) amine oxides of the formula $-\overset{\overset{\text{N}^+}{\mid}}{\text{O}^-}\text{R}_{30}\text{R}_{31}$ provided R₃₀ and R₃₁ are not hydrogen; or



wherein F and G are independently -OH, -SH, -OR₃₄, -SR₃₄ or -NR₃₂R₃₃; or

- (8) halogen, cyano, nitro, or azido. Alkyl, aryl and alkaryl groups on the substituents
- 10 of the above-defined alkyl groups may contain one additional substituent but are preferably unsubstituted. Examples of such radicals include, but are not limited to, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isoamyl, hexyl, octyl, nonyl, decyl, dodecyl, tetradecyl, hexadecyl, octadecyl and eicosyl. The term "alkenyl", alone or in combination, means an alkyl radical having
- 15 one or more double bonds. Examples of such alkenyl radicals include, but are not limited to, ethenyl, propenyl, 1-butenyl, cis-2-butenyl, trans-2-butenyl, isobutylenyl, cis-2-pentenyl, trans-2-pentenyl, 3-methyl-1-butenyl, 2,3-dimethyl-2-butenyl, 1-pentenyl, 1-hexenyl, 1-octenyl, decenyl, dodecenyl, tetradecenyl, hexadecenyl, cis- and trans- 9-octadecenyl, 1,3-pentadienyl, 2,4-pentadienyl, 2,3-
- 20 pentadienyl, 1,3-hexadienyl, 2,4-hexadienyl, 5,8,11,14-eicosatetraenyl, and 9,12,15-octadecatrienyl. The term "alkynyl", alone or in combination, means an alkyl radical having one or more triple bonds. Examples of such alkynyl groups include, but are not limited to, ethynyl, propynyl (propargyl), 1-butyne, 1-octynyl, 9-octadecynyl, 1,3-pentadiynyl, 2,4-pentadiynyl,
- 25 1,3-hexadiynyl, and 2,4-hexadiynyl. The term "cycloalkyl", alone or in combination means a cycloalkyl radical containing from 3 to about 10, preferably from 3 to about

-20-

- 8, and most preferably from 3 to about 6, carbon atoms. Examples of such cycloalkyl radicals include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, and perhydronaphthyl. The term "cycloalkylalkyl" means an alkyl radical as defined above which is substituted by a cycloalkyl radical as defined above. Examples of cycloalkylalkyl radicals include, but are not limited to, cyclohexylmethyl, cyclopentylmethyl, (4-isopropylcyclohexyl)methyl, (4-*t*-butyl-cyclohexyl)methyl, 3-cyclohexylpropyl, 2-cyclohexylmethylpentyl, 3-cyclopentyl-methylhexyl, 1-(4-neopentylcyclohexyl)methylhexyl, and 1-(4-isopropylcyclohexyl)-methylheptyl. The term "cycloalkylcycloalkyl" means a cycloalkyl radical as defined above which is substituted by another cycloalkyl radical as defined above. Examples of cycloalkylcycloalkyl radicals include, but are not limited to, cyclohexylcyclopentyl and cyclohexylcyclohexyl. The term "cycloalkenyl", alone or in combination, means a cycloalkyl radical having one or more double bonds. Examples of cycloalkenyl radicals include, but are not limited to, cyclopentenyl, cyclohexenyl, cyclooctenyl, cyclopentadienyl, cyclohexadienyl and cyclooctadienyl. The term "cycloalkenylalkyl" means an alkyl radical as defined above which is substituted by a cycloalkenyl radical as defined above. Examples of cycloalkenylalkyl radicals include, but are not limited to, 2-cyclohexen-1-ylmethyl, 1-cyclopenten-1-ylmethyl, 2-(1-cyclohexen-1-yl)ethyl, 3-(1-cyclopenten-1-yl)propyl, 1-(1-cyclohexen-1-ylmethyl)pentyl, 1-(1-cyclopenten-1-yl)hexyl, 6-(1-cyclohexen-1-yl)hexyl, 1-(1-cyclopenten-1-yl)nonyl and 1-(1-cyclohexen-1-yl)nonyl. The terms "alkylcycloalkyl" and "alkenylcycloalkyl" mean a cycloalkyl radical as defined above which is substituted by an alkyl or alkenyl radical as defined above. Examples of alkylcycloalkyl and alkenylcycloalkyl radicals include, but are not limited to, 2-ethylcyclobutyl, 1-methylcyclopentyl, 1-hexylcyclopentyl, 1-methyl-cyclohexyl, 1-(9-octadecenyl)cyclopentyl and 1-(9-octadecenyl)cyclohexyl. The terms "alkylcycloalkenyl" and "alkenylcycloalkenyl" means a cycloalkenyl radical as defined above which is substituted by an alkyl or alkenyl radical as defined above. Examples of alkylcycloalkenyl and alkenylcycloalkenyl radicals include, but are not limited to, 1-methyl-2-cyclopentyl, 1-hexyl-2-cyclopentenyl, 1-ethyl-2-cyclohexenyl, 1-butyl-2-cyclohexenyl, 1-(9-octadecenyl)-2-cyclohexenyl and 1-(2-pentenyl)-2-cyclohexenyl. The term "aryl", alone or in combination, means a phenyl or naphthyl radical which optionally carries one or more substituents selected from alkyl, cycloalkyl, cycloalkenyl, aryl, heterocycle, alkoxyaryl, alkaryl, alkoxy, halogen, hydroxy, amine, cyano, nitro, alkylthio, phenoxy, ether, trifluoromethyl and the like, such as phenyl, *p*-tolyl,

4-methoxyphenyl, 4-(tert-butoxy)phenyl, 4-fluorophenyl, 4-chlorophenyl, 4-hydroxyphenyl, 1-naphthyl, 2-naphthyl, and the like. The term "aralkyl", alone or in combination, means an alkyl or cycloalkyl radical as defined above in which one hydrogen atom is replaced by an aryl radical as defined above, such as benzyl, 2-phenylethyl, and the like. The term "heterocyclic" means ring structures containing at least one other kind of atom, in addition to carbon, in the ring. The most common of the other kinds of atoms include nitrogen, oxygen and sulfur. Examples of heterocyclics include, but are not limited to, pyrrolidinyl, piperidyl, imidazolidinyl, tetrahydrofuryl, tetrahydrothienyl, furyl, thienyl, pyridyl, quinolyl, isoquinolyl, pyridazinyl, pyrazinyl, indolyl, imidazolyl, oxazolyl, thiazolyl, pyrazolyl, pyridinyl, benzoxadiazolyl, benzothiadiazolyl, triazolyl and tetrazolyl groups. The term "saturated, partially saturated or unsaturated cyclic" means fused ring structures in which 2 carbons of the ring are also part of the fifteen-membered or sixteen membered macrocyclic ligand. The ring structure can contain 3 to 20 carbon atoms, preferably 5 to 10 carbon atoms, and can also contain one or more other kinds of atoms in addition to carbon. The most common of the other kinds of atoms include nitrogen, oxygen and sulfur. The ring structure can also contain more than one ring and can contain pendant substituents or functional groups as defined in the definition of alkyl groups herein. When the saturated, partially saturated or unsaturated cyclic is fused to the portion of the sixteen-membered ring having three carbons between nitrogens, the fused ring structure will contain 4 to 20 carbon atoms, preferably 5 to 10 carbon atoms, and can also contain one or more other kinds of atoms in addition to carbon. The term "saturated, partially saturated or unsaturated ring structure" means a ring structure in which one carbon of the ring is also part of the fifteen-membered or sixteen-membered macrocyclic ligand. The ring structure can contain 3 to 20, preferably 5 to 10, carbon atoms and can also contain nitrogen, oxygen and/or sulfur atoms. The term "nitrogen containing heterocycle" means ring structures in which 2 carbons and a nitrogen of the ring are also part of the fifteen-membered or sixteen-membered macrocyclic ligand. The ring structure can contain 2 to 20, preferably 4 to 10, carbon atoms, can be partially or fully unsaturated or saturated and can also contain nitrogen, oxygen and/or sulfur atoms in the portion of the ring which is not also part of the fifteen-membered macrocyclic ligand. The term "organic acid anion" refers to carboxylic acid anions having from about 1 to about 18 carbon atoms. The term "halide" means chloride or bromide.

As utilized herein, the term "radicals attached to the α -carbon of

α -amino acids" means "R" groups that are derived from the D or L forms of the amino acids Alanine, Aspartic acid, Arginine, Asparagine, Cysteine, Glycine, Glutamic acid, Glutamine, Histidine, Isoleucine, Leucine, Lysine, Methionine, Proline, Phenylalanine, Serine, Tryptophan, Threonine, Tyrosine, Valine and /or the

- 5 "R" groups of unnatural α -amino acids such as alkyl, ethyl, butyl, tert-butyl, cycloalkyl, phenyl, alkenyl, allyl, alkynyl, aryl, heteroaryl, polycycloalkyl, polycycloaryl, polycycloheteroaryl, imines, aminoalkyl, hydroxyalkyl, hydroxyl, phenol, amine oxides, thioalkyl, carboalkoxyalkyl, carboxylic acids and their derivatives, keto, ether, aldehyde, amine, nitrile, halo, thiol, sulfoxide, sulfone, 10 sulfonic acid, sulfide, disulfide, phosphonic acid, phosphinic acid, phosphine oxides, sulfonamides, amides, amino acids, peptides, proteins, carbohydrates, nucleic acids, fatty acids, lipids, nitro, hydroxylamines, hydroxamic acids, thiocarbonyls, borates, boranes, boraza, silyl, siloxy, silaza, and combinations thereof.

- The diamines of the invention can be prepared by any conventional method such as those described in Jung, S-H and Kohn, K., *J. Am. Chem. Soc.*, 15 1985, 107, 2931-43 and Roskamp, E.J. and Pederson, S.F., *J. Am. Chem. Soc.*, 1987, 109, 3152-4 or are commercially available. Examples of suitable diamines include, but are not limited to, *trans* 1,2-diaminocyclohexane, 2,3-diaminobutane, 2,2-dimethyl,1,3-diaminopropane, and 2,2-dimethyl,1,2-diaminoethane.

- 20 The triamines of the invention can be prepared by any conventional method such as by coupling an N-protected amino acid with a vicinal diamine, e.g. ethylene diamine, 1R,2R-diaminocyclohexane and the like, followed by removing the protecting group, and further optionally reducing the carbonyl group. The middle nitrogen of this reduced triamine can then be protected if desired according to the method described in John D. Prush, Laura A. Birchenough, & Melissa S. Egbertson, 25 *Synth. Comm.*, Vol. 22, pp. 2357-60 (1992). Examples of suitable triamines include, but are not limited to, 4-(p-toluenesulfonyl)-1,4,7-triazaheptane and *cis* -1-N-(*trans*-2'-aminocyclohexyl)-diaminocyclohexane.

- The dicarboxylic acids of the invention can be prepared by any 30 conventional method such as those set out in the Examples. Examples of suitable dicarboxylic acids include, but are not limited to, those used in the Examples, such as 3,6-bis(p-toluenesulfonyl)-3,6-diazaoctanedioic acid.

- The amount of diamine employed according to the invention can be conveniently expressed in terms of a molar ratio of diamine to dicarboxylic acid. 35 Broadly, the molar ratio of diamine to dicarboxylic acid will be about 0.5:1 to about 2:1, preferably about 0.8:1 to about 1.2:1, and most preferably about 1:1.

The amount of triamine employed according to the invention can be conveniently expressed in terms of a molar ratio of triamine to dicarboxylic acid. Broadly, the molar ratio of triamine to dicarboxylic acid will be about 0.5:1 to about 2:1, preferably about 0.8:1 to about 1.2:1, and most preferably about 1:1.

- 5 As used herein, the term "coupling agent" means an agent capable of effecting the dehydrative coupling of a carboxylic acid group of the dicarboxylic acid with a terminal amine group of the diamine or triamine to form an amide bond between the reactants. The agents promote or facilitate the dehydrative coupling by activating the carboxylic acid group. Descriptions of such coupling agents are
- 10 included in general text books of peptide chemistry, such as E. Schroder & K.L. Lubke, "The Peptides", Vol. 1, Academic Press, New York, NY, 1965, pp. 2-128; K.D. Kopple, "Peptides and Amino acids", W.A. Benjamin, Inc., New York, NY, 1966, pp. 33-51; and M. Bodanszky, "Principles of Peptide Synthesis", Springer-Verlag, 1984, pp. 16-58. The selection of an appropriate coupling agent is within the
- 15 skill in the art. Suitable coupling agents are selected from the group consisting of (1) phosphoryl azides (e.g. diphenylphosphoryl azide), (2) carbodiimides (e.g. N,N'-diisopropylcarbodiimide, N,N'-dicyclohexyl-carbodiimide and N-ethyl-N'-(γ -dimethylaminopropylcarbodiimide), (3) cyanamides (e.g. N,N-dibenzyl-cyanamide), (4) ketenimines, (5) isoxazolium salts (e.g. N-ethyl-5-phenyl-isoxazolium-3'-
- 20 sulfonate), (6) monocyclic nitrogen containing heterocyclic amides of aromatic character containing 1 to 4 nitrogens in the ring such as imidazolides, pyrazolides and 1,2,4-triazolides (e.g. N,N'-carbonyldiimidazole and N,N-carbonyldi-1,2,4-triazole), (7) alkoxylated acetylene (e.g. ethoxyacetylene), (8) reagents which form a mixed anhydride with the carboxyl moiety of the dicarboxylic acid (e.g.
- 25 ethylchloroformate, isobutylchloroformate, isobutyryl chloride, phosphorous trichloride, 1,2-phenylene phosphorochloridite, phosphorous oxychloride, diphenylchlorophosphate, N-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline, bis(2-oxo-3-oxazolidinyl)phosphonic chloride and triphenylphosphite-imidazole), (9) nitrogen containing heterocyclic compounds having a hydroxy group on one ring
- 30 nitrogen (e.g. N-hydroxyphthalimide, N-hydroxy-succinimide and 1-hydroxybenzotriazole) or carbonates thereof used in conjunction with a coupling agent from one of the other groups (e.g. 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride and 1-hydroxybenzotriazole, and dicyclohexylcarbodiimide and 1-hydroxybenzotriazole, (10) reagents which form an
- 35 active ester with the carboxyl moiety of the dicarboxylic acid (e.g. bromoacetonitrile,

-24-

substituted phenols such as p-nitrophenol, pentachlorophenol, pentafluorophenol, hydroxybenzotriazole, N-hydroxysuccinimide, thiophenol, thiophenol and 2-hydroxypyridine), (11) benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate, (12) re-dox based coupling agents (e.g. triphenylphosphine-2,2'-dipyridyldisulfide) and (13) mixtures thereof. The currently preferred coupling agents are diphenylphosphoryl azide, and 1-(3-dimethyl-aminopropyl)-3-ethylcarbodiimide hydrochloride with hydroxybenzotriazole.

The amount of coupling agent employed according to the invention can be conveniently expressed in terms of a molar ratio of coupling agent to dicarboxylic acid. Broadly, the molar ratio of coupling agent to dicarboxylic acid will be 2:1 to about 5:1, preferably 2:1 to about 3:1. If an anhydride of the dicarboxylic acid is used, the molar ratio of coupling agent to anhydride of the dicarboxylic acid will be 1:1 to about 3:1, preferably 1:1 to about 2:1.

Suitable bases for use in the invention are bases that are nonreactive with the coupling agent activated dicarboxylic acid and include trialkylamines, pyridines, imidazoles, and the like. Examples of suitable bases include, but are not limited to, triethylamine, pyridine, N-methylmorpholine and imidazole.

The amount of base employed according to the invention is the amount necessary to maintain the pH of the reaction medium to about 7 to about 9. The amount of base can be conveniently expressed in terms of a molar ratio of base to dicarboxylic acid. Broadly, the molar ratio of base to dicarboxylic acid will be 2:1 to about 4:1, preferably 2:1 to about 3:1. If the diamine or triamine is utilized as the diammonium salt which is converted *in situ* to the amine, the amount of base required will be increased to account for the base used to convert the diammonium salt to the amine.

Suitable solvents for use in the invention are non-reactive solvents in which the starting materials are at least partially soluble such as polar organic solvents. Examples of suitable solvents include, but are not limited to, dimethylformamide, dimethylacetamide, tetrahydrofuran, N-methyl-2-pyrrolidone, acetonitrile and mixtures thereof.

The amount of solvent employed according to the invention can be conveniently expressed in terms of molarity (moles/L or M). Broadly, the molarity will be about 0.001M to about 1 M, preferably about 0.005 M to about 0.1 M.


The temperature and time used in the process of the invention will depend on the particular reaction involved and will vary depending on factors such as coupling agent used, solvent used, concentration of the reaction mixture, and the solubility of the reactants. Based on the Examples herein, the temperature and time

-25-

for any particular combination of reactants, solvent, coupling agent, concentration, etc. will be readily apparent to those skilled in the art. For example, a reaction temperature is about -30 °C to about 25 °C will generally be acceptable for most reactions. However, when the ester of the dicarboxylic acid is an alkyl ester an elevated reaction temperature is required, i.e. the reaction temperature will be within 40°C of the boiling point of the solvent being used in the reaction. The time required will depend on the above factors as well as the temperature used.

The pentaazamacrocycles of the present invention can possess one or more asymmetric carbon atoms and are thus capable of existing in the form of optical isomers as well as in the form of racemic or nonracemic mixtures thereof. The optical isomers can be obtained by resolution of the racemic mixtures according to conventional processes, for example by formation of diastereoisomeric salts by treatment with an optically active acid. Examples of appropriate acids are tartaric, diacetyltartaric, dibenzoyltartaric, ditoluoyltartaric and camphorsulfonic acid and then separation of the mixture of diastereoisomers by crystallization followed by liberation of the optically active bases from these salts. A different process for separation of optical isomers involves the use of a chiral chromatography column optimally chosen to maximize the separation of the enantiomers. Still another available method involves synthesis of covalent diastereoisomeric molecules by reacting one or more secondary amine group(s) of the compounds of the invention with an optically pure acid in an activated form or an optically pure isocyanate. The synthesized diastereoisomers can be separated by conventional means such as chromatography, distillation, crystallization or sublimation, and then hydrolyzed to deliver the enantiomerically pure ligand. The optically active compounds of the invention can likewise be obtained by utilizing optically active starting materials, such as natural amino acids.

The commonly accepted mechanism of action of the manganese-based SOD enzymes involves the cycling of the manganese center between the two oxidation states (II,III). See J. V. Bannister, W. H. Bannister, and G. Rotilio, Crit. Rev. Biochem., 22, 111-180 (1987).



35

The formal redox potentials for the $O_2/O_2^{\cdot -}$ and HO_2^{\cdot}/H_2O_2 couples at pH = 7 are -0.33 v and 0.87 v, respectively. See A. E. G. Cass, in *Metalloproteins: Part 1, Metal Proteins with Redox Roles*, ed. P. Harrison, P. 121. Verlag Chemie (Weinheim, GDR) (1985). For the above disclosed mechanism, these potentials
5 require that a putative SOD catalyst be able to rapidly undergo oxidation state changes in the range of -0.33 v to 0.87 v.

The complexes derived from Mn(II) and the general class of C-substituted [15]aneN₅ ligands described herein have all been characterized using cyclic voltammetry to measure their redox potential. The C-substituted complexes
10 described herein have reversible oxidations of about +0.7 v (SHE). Coulometry shows that this oxidation is a one-electron process; namely it is the oxidation of the Mn(II) complex to the Mn(III) complex. Thus, for these complexes to function as SOD catalysts, the Mn(III) oxidation state is involved in the catalytic cycle. This means that the Mn(III) complexes of all these ligands are equally competent as SOD
15 catalysts, since it does not matter which form (Mn(II) or Mn(III)) is present when superoxide is present because superoxide will simply reduce Mn(III) to Mn(II) liberating oxygen.

The manganese complexes of the present invention can be utilized to treat numerous inflammatory disease states and disorders. For example, reperfusion
20 injury to an ischemic organ, e.g., reperfusion injury to the ischemic myocardium, surgically-induced ischemia, inflammatory bowel disease, rheumatoid arthritis, osteoarthritis, psoriasis, organ transplant rejections, radiation-induced injury, oxidant-induced tissue injuries and damage, atherosclerosis, thrombosis, platelet aggregation, stroke, acute pancreatitis, insulin-dependent diabetes mellitus,
25 disseminated intravascular coagulation, fatty embolism, adult and infantile respiratory distress, metastasis and carcinogenesis.

Activity of the compounds or complexes of the present invention for catalyzing the dismutation of superoxide can be demonstrated using the stopped-flow kinetic analysis technique as described in Riley, D.P., Rivers, W.J. and Weiss,
30 R.H., "Stopped-Flow Kinetic Analysis for Monitoring Superoxide Decay in Aqueous Systems," *Anal. Biochem.*, **196**, 344-349 (1991), which is incorporated by reference herein. Stopped-flow kinetic analysis is an accurate and direct method for quantitatively monitoring the decay rates of superoxide in water. The stopped-flow kinetic analysis is suitable for screening compounds for SOD activity and catalytic
35 activity of the compounds or complexes of the present invention for dismutating

-27-

superoxide, as shown by stopped-flow analysis, correlate to treating the above disease states and disorders.

Total daily dose administered to a host in single or divided doses may be in amounts, for example, from about 1 to about 100 mg/kg body weight daily and more usually about 3 to 30 mg/kg. Unit dosage compositions may contain such amounts of submultiples thereof to make up the daily dose.

The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration.

The dosage regimen for treating a disease condition with the compounds and/or compositions of this invention is selected in accordance with a variety of factors, including the type, age, weight, sex, diet and medical condition of the patient, the severity of the disease, the route of administration, pharmacological considerations such as the activity, efficacy, pharmacokinetic and toxicology profiles of the particular compound employed, whether a drug delivery system is utilized and whether the compound is administered as part of a drug combination. Thus, the dosage regimen actually employed may vary widely and therefore may deviate from the preferred dosage regimen set forth above.

The compounds of the present invention may be administered orally, parenterally, by inhalation spray, rectally, or topically in dosage unit formulations containing conventional nontoxic pharmaceutically acceptable carriers, adjuvants, and vehicles as desired. Topical administration may also involve the use of transdermal administration such as transdermal patches or iontophoresis devices. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection, or infusion techniques.

Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

Suppositories for rectal administration of the drug can be prepared by mixing the drug with a suitable nonirritating excipient such as cocoa butter and

polyethylene glycols which are solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum and release the drug.

Solid dosage forms for oral administration may include capsules, tablets, pills, powders, granules and gels. In such solid dosage forms, the active compound may be admixed with at least one inert diluent such as sucrose lactose or starch. Such dosage forms may also comprise, as in normal practice, additional substances other than inert diluents, e.g., lubricating agents such as magnesium stearate. In the case of capsules, tablets, and pills, the dosage forms may also comprise buffering agents. Tablets and pills can additionally be prepared with enteric coatings.

Liquid dosage forms for oral administration may include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs containing inert diluents commonly used in the art, such as water. Such compositions may also comprise adjuvants, such as wetting agents, emulsifying and suspending agents, and sweetening, flavoring, and perfuming agents.

While the compounds of the invention can be administered as the sole active pharmaceutical agent, they can also be used in combination with one or more compounds which are known to be effective against the specific disease state that one is targeting for treatment.

The compounds or complexes of the invention can also be utilized as MRI contrast agents. A discussion of the use of contrast agents in MRI can be found in patent application Serial No. 08/397,469, which is incorporated by reference herein.

Contemplated equivalents of the general formulas set forth above for the compounds and derivatives as well as the intermediates are compounds otherwise corresponding thereto and having the same general properties such as tautomers of the compounds and such as wherein one or more of the various R groups are simple variations of the substituents as defined therein, e.g., wherein R is a higher alkyl group than that indicated, or wherein the tosyl groups are other nitrogen or oxygen protecting groups or wherein the O-tosyl is a halide. Anions having a charge other than 1, e.g., carbonate, phosphate, and hydrogen phosphate, can be used instead of anions having a charge of 1, so long as they do not adversely affect the overall activity of the complex. However, using anions having a charge other than 1 will result in a slight modification of the general formula for the complex set forth above. In addition, where a substituent is designated as, or can be, a hydrogen, the exact chemical nature of a substituent which is other than hydrogen at that position, e.g., a hydrocarbyl radical or a halogen, hydroxy, amino and the like functional group, is not

-29-

critical so long as it does not adversely affect the overall activity and/or synthesis procedure. Further, it is contemplated that manganese(III) complexes will be equivalent to the subject manganese(II) complexes.

5 The chemical reactions described above are generally disclosed in terms of their broadest application to the preparation of the compounds of this invention. Occasionally, the reactions may not be applicable as described to each compound included within the disclosed scope. The compounds for which this occurs will be readily recognized by those skilled in the art. In all such cases, either the reactions can be successfully performed by conventional modifications known to those skilled
10 in the art, e.g., by appropriate protection of interfering groups, by changing to alternative conventional reagents, by routine modification of reaction conditions, and the like, or other reactions disclosed herein or otherwise conventional, will be applicable to the preparation of the corresponding compounds of this invention. In all preparative methods, all starting materials are known or readily preparable from
15 known starting materials.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way
20 whatsoever.

EXAMPLES

All reagents were used as received without purification unless otherwise indicated. All NMR spectra were obtained on a Varian VXR-300 or VXR-400 nuclear magnetic resonance spectrometer. Qualitative and quantitative mass
25 spectroscopy was run on a Finnigan MAT90, a Finnigan 4500 and a VG40-250T using m-nitrobenzyl alcohol(NBA) or m-nitrobenzyl alcohol/LiCl (NBA+Li); or dithiothreitol/dithioerythritol (DTT/DTE), under fast atom bombardment (FAB) or chemical ionization (CI) conditions. Melting points (mp) are uncorrected.

The following abbreviations relating to amino acids and their protective
30 groups are in accordance with the recommendation by IUPAC-IUB Commission on Biochemical Nomenclature (Biochemistry, 11, 1726 (1972)) and common usage.

-30-

	TFA	Trifluoroacetate
	DMF	Dimethylformamide
	HOBT·H ₂ O	1-Hydroxy-(1 <i>H</i>)-benzotriazole monohydrate
5	EDC·HCl	1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride
	TEA	Triethylamine
	DMSO	Dimethylsulfoxide
	THF	Tetrahydrofuran
10	DPPA	Diphenylphosphoryl azide
	DMPU	Dimethylpropyleneurea
	c	concentration, g/cc
	DME	1,2-Dimethoxyethane

15

Example 11A. Synthesis of *trans*-5,6-Dimethyl-3,8-dioxo-1,10,13-tris(p-toluenesulfonyl)-1,4,7,10,13-pentaazacyclopentadecane

- A solution of D,L-2,3-diaminobutane dihydrochloride containing 20% of the meso isomer (8.05 g, 50.0 mmole), 3,6,9-tris(p-toluenesulfonyl)-3,6,9-triazaundecanedioic acid prepared as in EP Patent Application 0 524 161 A1 (34.1 g, 50.0 mmole) and triethylamine (20.2 g, 200 mmole) in degassed anhydrous DMF (10.0 L) under a dry argon atmosphere was cooled to -45 to -50° C. Diphenylphosphorylazide (33.0 g, 120 mmole) was then added dropwise over 5 minutes and the resulting solution was allowed to warm to room temperature while stirring for 14 h. The solvent was then removed *in vacuo* and residual DMF was removed by co-evaporation with H₂O (2 x 1.0 L). The residue was dissolved in acetonitrile (300 ml) and methanol (700 ml) was added in portions as the product crystallized. The solid was filtered, washed with methanol and dried to give 17.2 g of colorless needles. The filtrate was concentrated to an oil and residual DMF was removed by co-evaporation with H₂O *in vacuo*. The residue was dissolved in a mixture of ethyl acetate (500 ml) and H₂O (500 ml) and the layers were separated. The ethyl acetate was washed with 0.1 N HCl (2 x 200 ml), saturated NaHCO₃ solution (2 x 200 ml) and saturated NaCl solution (500 ml), at which point a large mass of crystals formed in the ethyl acetate layer. The mixture was filtered and the crystals were washed with ethyl acetate and dried to give an additional 3.4 g of the

-31-

product as colorless needles. Recrystallization from CHCl_3 - methanol gave 21.7 g (59% yield) of the racemic product as colorless needles: mp 250 - 253 °C dec.; ^1H NMR (CDCl_3 , 400 MHz) δ 1.21 (d, J = 5.6 Hz, 6H), 2.45 (s, 9H), 3.20 (m, 4H), 3.39 (m, 2H), 3.46 (d, J = 16.4 Hz, 2H), 3.55 (m, 2H), 3.89 (m, 2H), 3.91 (d, J = 16.4 Hz, 2H), 6.55 (d, J = 6.2 Hz, 2H), 7.35 (m, 6H), 7.71 (m, 6H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 18.51, 21.58, 49.80, 50.70, 51.61, 54.49, 127.55, 127.68, 129.97, 130.09, 133.97, 134.33, 143.96, 144.47, 168.20; FAB mass spectrum (NBA+Li) m/z 740 $[\text{M} + \text{H}]^+$.

10 1B. Synthesis of *trans*-2,3-Dimethyl-1,4,7,10,13-pentaazacyclopentadecane

- To a stirred suspension of *trans*-5,6-dimethyl-1,10,13-tris(*p*-toluenesulfonyl)-1,4,7,10,13-pentaazacyclopentadecane-3,8-dione prepared as in example 1A (21.6 g, 29.4 mmole) in anhydrous THF (590 ml) under a dry argon atmosphere was added a solution of 1.0 M LiAlH_4 in THF (368 ml, 368 mmole) dropwise over 5 minutes.
- 15 The resulting clear yellow solution was refluxed for 32 h (by which time it had become heterogeneous) and was then cooled to 0 °C. The mixture was quenched by the dropwise addition of 10% Na_2SO_4 (18 ml) while cooling in an ice bath. The mixture was then filtered under argon and the solids were refluxed for 30 minutes with anhydrous THF (1.0 L). Following filtration, the solids were refluxed with
- 20 THF (1.0 L) and the mixture was filtered again. The filtrates were evaporated *in vacuo* and redissolved in THF, combined and filtered. The solvent was removed *in vacuo* and the residue was dissolved in hot acetonitrile (50 ml). The cloudy solution was filtered and the filtrate was concentrated to a volume of 20 ml. Crystallization was initiated by briefly touching a piece of dry ice to the outside of the flask. As
- 25 crystallization proceeded, the flask was slowly cooled to -20 °C. Following filtration, the crystals were purified by recrystallization from acetonitrile to give 1.87 g (26.1% yield) of the product as colorless needles: mp (under argon) 81 - 83 °C; ^1H NMR (C_6D_6 , 300 MHz) δ 0.88 (m, 6H), 1.46 (br s, 5H), 2.10 (m, 2H), 2.34 (m, 2H), 2.52 (m, 10H), 2.64 (m, 2H), 2.76 (m, 2H); ^{13}C NMR (C_6D_6 , 100 MHz) δ 17.31,
- 30 47.29, 48.47, 48.87, 49.34, 59.46; CI mass spectrum (CH_4) m/z 244 $[\text{M} + \text{H}]^+$.

1C. Synthesis of [Manganese(II) dichloro (D,L-2,3-Dimethyl-1,4,7,10,13-pentaazacyclopentadecane)]

-32-

To a stirred solution of anhydrous MnCl_2 (383 mg, 3.04 mmol) in anhydrous methanol was added D,L-2,3-dimethyl-1,4,7,10,13-pentaazacyclopentadecane prepared as in example 1B (740 mg, 3.04 mmol). The solution was refluxed for 1h, and then the solvent was removed *in vacuo*. The residue was dissolved in hot THF
5 and filtered through Celite™. The filtrate was concentrated *in vacuo* to a volume of 5 ml. Ethyl ether was added to the warm solution resulting in the precipitation of a white solid which was collected by filtration. After drying *in vacuo*, a yield of 0.81 g (72%) was obtained; FAB mass spectrum (NBA) m/z (relative intensity) 333/335 (100/30) $[\text{M} - \text{Cl}]^+$; Anal. Calcd. for $\text{C}_{12}\text{H}_{29}\text{N}_5\text{Cl}_2\text{Mn}$: C, 39.03; H, 7.92; N, 18.97;
10 Cl, 19.20. Found: C, 38.95; H, 7.96; N, 18.88; Cl, 19.25.

Example 2

2A. Preparation of Dimethyl 2S,10S-bis(tert-butylthiomethyl)-4,8-dioxo-6-p-toluenesulfonyl-3,6,9-triazaundecanoate.

N-p-Toluenesulfonyliminodiacetic acid (7.73 g, 26.9 mmol), prepared according to example 5A, was placed in a 500 ml round bottomed flask equipped with a magnetic stirbar and under argon atmosphere. DMF (300 ml) was added to the flask, and the flask was cooled in an ice-water bath. HOBT·H₂O (8.25 g, 53.9
20 mmol) was added followed by EDC·HCl (12.38 g, 64.6 mmol). After 5 min, S-t-butyl-L-cysteine methyl ester hydrochloride (prepared from S-t-butyl-L-cysteine following: Pastuszak, J. J., Chimiak, A. *J. Org. Chem.* 1981, 46, 1868 - 1873) was added to the solution followed by triethylamine (7.51 ml, 53.9 mmol). The mixture turned a light orange color and precipitate began to form. The mixture was stirred and
25 allowed to slowly warm to room temperature. After ~16 h, the mixture was concentrated to ~75 ml volume *in vacuo*. Ethyl acetate (EtOAc) (300 ml) and aq. NaHCO_3 (200 ml) were added to the slurry. The layers were separated and the aqueous layer was extracted twice with 100 ml of EtOAc. The EtOAc layers were combined and washed with 0.5 N NaHSO_4 (200 ml), sat. NaHCO_3 (100 ml), 2 x
30 100 ml water, brine (100 ml), dried over MgSO_4 , filtered and concentrated to a pale yellow foam, 15.0 g (88%). ^1H NMR (CDCl_3 , 300 MHz) δ 1.28 (s, 18H), 2.40 (s, 3H), 2.86 - 3.01 (m, 4H), 3.73 (s, 6H), 3.90 (s, 4H), 4.73 (dd, $J = 5.7, 13.2$ Hz, 2H), 7.29 (d, $J = 8.1$ Hz, 2H), 7.39 (d, $J = 7.5$ Hz, 2H), 7.75 (d, $J = 8.4$ Hz, 2H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 21.6, 31.0, 30.8, 42.7, 52.1, 52.5, 52.6, 127.9,
35 129.8, 134.7, 144.3, 167.8, 170.8.

2B. Preparation of 2S,10S-Bis(tert-butylthiomethyl)-4,8-dioxo-6-p-toluenesulfonyl-3,6,9-triazaundecanoic acid

- NaOH (1.84 g, 46.0 mmol) in 50 ml of water was slowly added to a cooled
 5 solution of dimethyl 2S,10S-bis(tert-butylthiomethyl)-4,8-dioxo-6-p-toluenesulfonyl-3,6,9-triazaundecanoate (14.68 g, 23.16 mmol) in 100 ml of THF. After 1.5 h, aq. HCl (23.2 mmol) was added to the cooled solution and the THF was removed *in vacuo*. The mixture was then neutralized with the addition of 23 ml of 1N HCl. EtOAc was added and the pH of the aqueous layer was monitored. Additional 1N
 10 HCl was added to bring the aqueous layer to pH = 2.5 and the layers were separated. The aqueous layer was extracted twice with 100 ml of EtOAc. The EtOAc layers were combined, washed with 50 ml of water and 100 ml of brine, dried over MgSO₄, filtered and concentrated to a dry foam, 13.7 g (98%). ¹H NMR (DMSO-d₆, 300 MHz) δ 1.24 (s, 18H), 2.37 (s, 3H), 2.68 - 2.87 (m, 4H), 3.94 (s, 4H), 4.34 (dd, J
 15 = 7.1, 13.7 Hz, 2H), 7.36 (d, J = 8.1 Hz, 2H), 7.68 (d, J = 8.1 Hz, 2H), 8.69 (d, J = 8.1 Hz, 2H), 12.87 (br s, 2H). ¹³C NMR (DMSO-d₆, 100 MHz) δ 21.0, 29.6, 30.6, 42.2, 50.6, 52.8, 127.2, 129.5, 135.6, 143.2, 168.0, 171.4.

2C. Preparation of 6S,14S-Bis(tert-butylthiomethyl)-2R,3R-cyclohexano-5,8,12,15-tetraoxo-10-p-toluenesulfonyl-1,4,7,10,13-pentaazacyclopentadecane

- HOBT·H₂O (6.88 g, 44.9 mmol) and EDC·HCl (10.33 g, 53.9 mmol) were added to a cooled solution (0°C) of 2S,10S-bis(tert-butylthiomethyl)-4,8-dioxo-6-p-toluenesulfonyl-3,6,9-triazaundecanoic acid (13.60 g, 22.4 mmol) in 4 L of DMF. After 15 min, 1R,2R-diaminocyclohexane in 30 ml of DMF was added to the cooled
 25 solution. The solution was stirred under argon atmosphere and allowed to slowly warm to room temperature. After ~16 h, the solution was concentrated *in vacuo* to a slurry and EtOAc and aq. NaHCO₃ (~100 ml) were added. The layers were combined and separated. The aqueous layer was extracted with EtOAc, then all EtOAc layers were combined, washed with 0.5 N NaHSO₄, saturated NaHCO₃,
 30 water (2 times), brine, dried over MgSO₄, filtered and concentrated to yield crude solid. Pure product was isolated by trituration with EtOAc, as a white solid, weight 8.6 g (56%). ¹H NMR (300 MHz, DMSO-d₆) δ 1.00 - 1.39 (m, 22H), 1.52 - 1.69 (m, 2H), 1.69 - 1.88 (m, 2H), 2.39 (s, 3H), 2.50 - 2.75 (m, 4H), 3.32 - 3.52 (m, 2H), 3.66 (d, J = 17.3 Hz, 2H), 3.86 (d, J = 17.1 Hz, 2H), 4.09 - 4.24 (m, 2H),
 35 7.44 (d, J = 8.3 Hz, 2H), 7.76 (d, J = 8.1 Hz, 2H), 7.79 - 7.90 (m, 2H), 8.59 (d, J

-34-

= 7.1 Hz, 2H). ^{13}C NMR (100 MHz, DMSO-d_6) δ 21.0, 24.4, 29.0, 30.7, 31.9, 42.0, 52.1, 52.7, 54.1, 127.4, 129.9, 134.8, 143.8, 168.2, 168.8; FAB mass spectrum (NBA+Li) m/z 690 $[\text{M} + \text{Li}]^+$, 696 $[\text{M} - \text{H} + 2\text{Li}]^+$.

5 2D. Preparation of 2S,9S-Bis(tert-butylthiomethyl)-5R,6R-cyclohexano-1,4,7,10,13-pentaazacyclopentadecane

- A 500 ml round bottomed flask equipped with a magnetic stirbar and condenser under argon atmosphere was charged with 6S,14S-bis(tert-butylthiomethyl)-2R,3R-cyclohexano-5,8,12,15-tetraoxo-10-p-toluenesulfonyl-
- 10 1,4,7,10,13-pentaazacyclopentadecane (2.88 g, 4.21 mmol) in 100 ml of THF. LiAlH_4 (105 ml, 0.5M in DME, 52.5 mmol) was added to the mixture which was then heated to reflux. After ~16 h, the reaction was allowed to cool to room temperature then was cooled in an ice-water bath and quenched by the addition of 2.0 ml of water, 2.0 ml of 15% NaOH (aq.), and 6.0 ml of water. THF was added to the
- 15 mixture and the suspension was filtered under inert atmosphere several times. The filtrates were concentrated and the residue crystallized from acetonitrile to give 0.60 g of product (30%). ^1H NMR (CDCl_3 , 300 MHz) δ 0.88 - 1.08 (m, 2H), 1.15 - 1.30 (m, 2H), 1.29 (s, 18H), 1.64 - 1.79 (m, 2H), 1.94 - 2.25 (m, 11H), 2.45 - 2.89 (m, 12H), 2.90 - 3.05 (m, 2H), 3.06 - 3.18 (m, 2H); ^{13}C (CDCl_3 , 75 MHz) δ 24.9,
- 20 30.9, 31.2, 31.7, 42.0, 45.8, 48.5, 51.6, 58.0, 63.2; Exact mass ($\text{M} + \text{H}$) $^+$: calcd, 474.3664; found, 474.3686 ($\text{C}_{24}\text{H}_{52}\text{N}_5\text{S}_2$).

2E. Preparation of [Manganese(II) dichloro (2S,9S-t-butylthiomethyl-5R,6R-cyclohexano-1,4,7,10,13-pentaazacyclopentadecane)]

- 25 2S,9S-Bis(tert-butylthiomethyl)-5R,6R-cyclohexano-1,4,7,10,13-pentaazacyclopentadecane (269 mg, 0.568 mmol) was added to a hot methanol (MeOH) solution containing 90 mg of MnCl_2 . The solution was stirred overnight and then was evaporated to dryness. The resultant oil was dissolved in hot THF, filtered through CeliteTM, and evaporated to dryness to yield 275 mg (81%). FAB mass
- 30 spectrum m/z 563 $[\text{M} - \text{Cl}]^+$. Catalytic rate constant for the dismutation of superoxide at pH 8.1 and 21 °C, $k_{\text{cat}} = 1.29 \times 10^7 \text{ mol}^{-1}\text{sec}^{-1}$.

Example 33A. Synthesis of Ethyl 2R,3R-cyclohexano-1,4-diaza-5R-methyl-1-(p-toluenesulfonyl)-6-hexanoate

- 5 To a stirred solution of N-p-toluenesulfonyl-1R,2R-diaminocyclohexane, prepared as in U.S. Patent Application Serial No. 08/397,469, (9.98 g, 37.2 mmole) in anhydrous methylene chloride (CH_2Cl_2) (250 ml) under a dry argon atmosphere was added triethylamine (3.76 g, 37.2 mmole) and the resulting solution was cooled to 0° C. A solution of ethyl L-lactate trifluoromethanesulfonate (9.30 g, 37.2 mmole)
- 10 in anhydrous CH_2Cl_2 (50 ml) was added dropwise to the solution over 45 minutes, maintaining the temperature at 0° C. The mixture was then stirred for 16 h while allowing it to warm to room temperature. CH_2Cl_2 (500 ml) was added and the solution was washed with saturated NaHCO_3 solution (500 ml), H_2O (500 ml) and saturated NaCl solution (250 ml) and was dried (MgSO_4). After filtration, the solvent
- 15 was removed *in vacuo* to give 14.3 g of crude product as a yellow oil: ^1H NMR (CDCl_3 , 300 MHz) δ 0.94 (m, 1H), 1.10 - 1.34 (m, 9H), 1.51 - 1.68 (m, 3H), 1.86 (m, 1H), 2.20 (m, 2H), 2.44 (s, 3H), 2.52 (m, 1H), 3.29 (q, J = 6.8 Hz, 1H), 4.15 (q, J = 7.1 Hz, 2H), 5.30 (br s, 1H), 7.31 (d, J = 8.1 Hz, 2H), 7.76 (d, J = 8.3 Hz, 2H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 14.17, 19.76, 21.53, 24.10, 24.73, 32.24,
- 20 32.34, 54.82, 58.05, 59.71, 60.95, 127.28, 129.61, 137.10, 143.22, 176.10; FAB mass spectrum (DTT/DTE) m/z 369 $[\text{M} + \text{H}]^+$.

3B. Synthesis of Ethyl 4-benzoyloxycarbonyl-2R,3R-cyclohexano-1,4-diaza-5R-methyl-1-(p-toluenesulfonyl)-6-hexanoate

- 25 To a stirred solution of ethyl 2R,3R-cyclohexano-1,4-diaza-5R-methyl-1-(p-toluenesulfonyl)-6-hexanoate prepared as in Example 3A (14.2 g of crude product) in THF (250 ml) was added 2N NaOH (20 ml, 40.0 mmole) slowly at 0° C. Benzyl chloroformate (6.94 g, 40.7 mmole) was added in portions and the mixture was allowed to warm to room temperature. At 2 h, additional benzyl chloroformate (2.00
- 30 g, 11.7 mmole) was added, the pH was adjusted to 8 - 9 with 2N NaOH and the mixture was stirred for 18 h. After filtration, the solvent was removed *in vacuo* to give an oil which rapidly crystallized. The crude product was dissolved in a mixture of ethyl acetate (500 ml) and H_2O (500 ml) and the layers were separated. The ethyl acetate layer was washed with H_2O (2 x 500 ml), saturated NaCl solution and was
- 35 dried (MgSO_4). The solvent was removed *in vacuo* to give 26.7 g of a colorless oil.

-36-

Following trituration of the oil with hexanes, the product crystallized into a solid mass. Recrystallization from CH_2Cl_2 - hexanes gave 16.8 g (90.2% yield) of the product as colorless prisms: mp 119 - 121 °C; ^1H NMR ($\text{DMSO}-d_6$, 100 °C, 400 MHz) δ 0.98 - 1.18 (m, 6H), 1.40 (d, J = 7.0 Hz, 3H), 1.48 (br d, J = 12.3 Hz, 1H), 1.59 (br d, J = 11.0 Hz, 1H), 1.70 (br s, 1H), 1.86 (br s, 1H), 2.33 (s, 3H), 3.14 (br q, J = 7.3 Hz, 1H), 3.63 (br s, 1H), 3.92 - 4.01 (m, 3H), 4.87 - 4.97 (m, 2H), 6.88 (d, J = 7.2 Hz, 1H), 7.28 - 7.35 (m, 7H), 7.61 (d, J = 8.1 Hz, 2H); ^{13}C NMR ($\text{DMSO}-d_6$, 100° C, 100 MHz) δ 14.33, 16.50, 21.35, 24.82, 25.53, 29.78, 33.71, 53.87, 58.35, 61.05, 65.63, 66.76, 126.83, 127.91, 128.09, 128.65, 129.74, 137.06, 140.32, 142.56, 154.50, 172.02; FAB mass spectrum (NBA+Li) m/z 509 $[\text{M} + \text{H}]^+$.

3C. Synthesis of 4-Benzoyloxycarbonyl-2R,3R-cyclohexano-1,4-diaza-5R-methyl-1-(p-toluenesulfonyl)-6-hexanoic acid

To a stirred suspension of ethyl 4-benzoyloxycarbonyl-2R,3R-cyclohexano-1,4-diaza-5R-methyl-1-(p-toluenesulfonyl)-6-hexanoate prepared as in Example 3B (11.7 g, 23.2 mmole) was slowly added 2N NaOH (17.4 ml, 34.8 mmole) and the resulting clear solution was stirred for 3 h at room temperature and then at 50 °C for another 2 h. The methanol was removed *in vacuo*, H_2O (250 ml) was added and the aqueous solution was washed with ethyl ether (500 ml). The pH was then adjusted to 2 with 1N HCl and the product was extracted into ethyl acetate (3 x 500 ml). The extracts were combined, washed with H_2O (250 ml), saturated NaCl solution (250 ml) and were dried (MgSO_4). The solvent was removed *in vacuo* to give 10.76 g (97.8% yield) of the product as a white solid: ^1H NMR ($\text{DMSO}-d_6$) shows a doubling of peaks at 25° C due to conformational isomers; ^1H NMR ($\text{DMSO}-d_6$, 100° C, 300 MHz) δ 1.02 - 1.20 (m, 3H), 1.39 (d, J = 6.9 Hz, 3H), 1.39 - 1.61 (m, 2H), 1.89 (m, 3H), 2.34 (s, 3H), 3.17 (m, 1H), 3.56 (m, 1H), 3.97 (q, J = 6.9 Hz, 1H), 4.90 (m, 2H), 7.03 (br s, 1H), 7.27 - 7.37 (m, 7H), 7.61 (d, J = 7.9 Hz, 2H), 12.33 (br s, 1H); ^{13}C NMR ($\text{DMSO}-d_6$, 100° C, 100 MHz) δ 15.99, 20.79, 24.14, 24.95, 29.51, 33.18, 53.38, 56.86, 63.97, 66.19, 126.35, 127.14, 127.43, 128.10, 129.18, 136.53, 139.34, 142.02, 154.09, 173.55; FAB mass spectrum (NBA - HCl) m/z 475 $[\text{M} + \text{H}]^+$.

-37-

3D. Synthesis of Methyl 4-benzyloxycarbonyl-2R,3R-cyclohexano-5R,8R-dimethyl-6-oxo-1-(p-toluenesulfonyl)-1,4,7-triazanonanoate

To a stirred solution of 4-benzyloxycarbonyl-2R,3R-cyclohexano-1,4-diaza-5R-methyl-1-(p-toluenesulfonyl)-6-hexanoic acid prepared as in example 3C (10.6 g, 22.6 mmole), 1-hydroxybenzotriazole monohydrate (3.66 g, 23.9 mmole) and triethylamine (5.25 g, 51.9 mmole) in anhydrous DMF (250 ml) under a dry argon atmosphere was added 1-ethyl-3-(3-dimethyl-aminopropyl)carbodiimide hydrochloride (5.19 g, 27.1 mmole) and the mixture was stirred at room temperature for 30 minutes. Then, D-alanine methyl ester hydrochloride (3.47 g, 24.83 mmole) was added and the mixture was stirred at room temperature for 17 h. The solvent was then removed *in vacuo* and the residue was dissolved in a mixture of ethyl acetate (500 ml) and H₂O (500 ml). The layers were separated and the ethyl acetate layer was washed with 0.1 N HCl (2 x 250 ml), saturated NaHCO₃ solution (2 x 250 ml), saturated NaCl solution (250 ml) and was dried (MgSO₄). The solvent was removed *in vacuo* to give 11.8 g (93.4% yield) of a pale yellow solid: ¹H NMR (DMSO-d₆, 100° C, 400 MHz) δ 1.02 - 1.94 (m, 14H), 2.31 (s, 3H), 3.05 (m, 1H), 3.22 (m, 1H), 3.64 (s, 3H), 4.33 - 4.82 (m, 4H), 7.22 - 7.36 (m, 7H), 7.58 (d, J = 7.5 Hz, 2H), 7.82 (br s, 1H), 8.13 (br s, 1H); ¹³C NMR (DMSO-d₆, 100 MHz) δ 16.15, 16.75, 20.78, 24.07, 24.96, 30.62, 34.43, 47.88, 51.78, 53.46, 54.80, 59.39, 66.24, 126.37, 127.21, 127.58, 128.20, 129.18, 136.51, 139.39, 141.90, 154.70, 172.38, 173.06; FAB mass spectrum (NBA - Li) m/z 566 [M + Li]⁺.

3E. Synthesis of Dimethyl 6-benzyloxycarbonyl-4R,5R-cyclohexano-7R,10R-dimethyl-8-oxo-3-(p-toluenesulfonyl)-3,6,9-triazaundecanedioate

To a stirred solution of methyl 4-benzyloxycarbonyl-2R,3R-cyclohexano-5R,8R-dimethyl-6-oxo-1-(p-toluenesulfonyl)-1,4,7-triazanonanoate prepared as in example 3D (10.2 g, 18.2 mmole) in anhydrous DMF (100 ml) under a dry argon atmosphere was added methyl bromoacetate (5.58 g, 36.5 mmole) and the solution was cooled to 0° C. Sodium hydride (547 mg, 80% in oil, 18.2 mmole) was then added and mixture was allowed to warm to room temperature while stirring. At 2 h, additional NaH (547 mg, 80% in oil, 18.2 mmole) and methyl bromoacetate (5.58 g, 36.5 mmole) were added at 0° C and the mixture was stirred at 0° C. At 5.25 h, more NaH (274 mg, 80% in oil, 9.12 mmole) and methyl bromoacetate (2.79 g, 18.2 mmole) were added at 0° C and the mixture was stirred for another 30 minutes while warming to room temperature. The mixture was then quenched by pouring into H₂O

-38-

- (500 ml). The solvent was then removed *in vacuo* and the residue was dissolved in a mixture of ethyl acetate (1 L) and H₂O (1 L). The layers were separated and the ethyl acetate layer was washed with H₂O (2 X 1 L) and saturated NaCl solution (500 ml) and was dried (MgSO₄). The solvent was removed *in vacuo* to give 12.4 g of a pale yellow solid: FAB mass spectrum (NBA - Li) *m/z* 638 [M + Li]⁺. This crude product was not purified and was hydrolyzed in Example 3F.

3F. Synthesis of 6-Benzoyloxycarbonyl-4R,5R-cyclohexano-7R,10R-dimethyl-8-oxo-3-(p-toluenesulfonyl)-3,6,9-triazaundecanedioic acid

- To a solution of dimethyl 6-benzoyloxycarbonyl-4R,5R-cyclohexano-7R,10R-dimethyl-8-oxo-3-(p-toluenesulfonyl)-3,6,9-triazaundecanedioate prepared as in Example 3E (12.3 g, 19.5 mmole) in anhydrous methanol (250 ml) was slowly added a solution of 2N sodium hydroxide (30 ml, 60.0 mmole) and the bright yellow solution was stirred at room temperature for 2 h and then at 50° C for 2 h. Then, 10 ml of 2 N sodium hydroxide was added and the solution was stirred at 50° C for 2.5 h at which time the reaction was complete. The solvent was removed *in vacuo* and the residue was dissolved in H₂O (500 ml). The aqueous solution was washed with ethyl ether (2 x 500 ml) and the pH was adjusted to 1 with 1 N HCl. After saturating the aqueous mixture with NaCl, the product was extracted into ethyl acetate (3 x 500 ml), the extracts were combined, dried over MgSO₄, and the solvent was removed *in vacuo*. Crystallization of the crude product from ethyl acetate - ethyl ether resulted in the formation of a gelatinous precipitate. The solid was filtered, washed with ethyl ether and dried to give 5.98 g (50.7 % for two steps) of the product as a white solid: mp 155 °C; ¹H NMR (DMSO-d₆, 100° C, 400 MHz) δ 0.90 - 2.10 (m, 14 H), 2.36 (s, 3H), 3.70 - 4.35 (m, 6H), 5.03 (m, 2H), 7.29 - 7.35 (m, 7H), 7.69 (d, J = 7.3 Hz, 2H), 8.00 (br s, (1H), 12.20 (br s, 2H); ¹³C NMR (DMSO-d₆, 100° C, 100 MHz) δ 17.26, 21.37, 21.46, 25.14, 25.39, 30.17, 31.40, 46.20, 49.15, 58.91, 59.70, 67.00, 127.95, 128.05, 128.47, 129.73, 129.84, 137.15, 138.12, 143.56, 155.53, 171.22, 171.31, 173.80; FAB mass spectrum (NBA - Li) *m/z* (relative intensity) 610 (74) [M + Li]⁺, 616 (100) [M + 2Li - H]⁺, 622 (87) [M + 3Li - 2H]⁺.

3G. Synthesis of 4-Benzoyloxycarbonyl-2R,3R,11R,12R-bis(cyclohexano)-5R,8R-dimethyl-6,9,14-trioxo-1,4,7,10,13-pentaaza-1-p-toluenesulfonyl-cyclopentadecane

- To a stirred solution of 6-benzoyloxycarbonyl-4R,5R-cyclohexano-7R,10R-dimethyl-8-oxo-3-(p-toluenesulfonyl)-3,6,9-triazaundecanedioic acid prepared as in

-39-

Example 3F (5.93 g, 9.83 mmole) and 1R,2R-diaminocyclohexane (1.12 g, 9.8 mmole) in degassed, anhydrous DMF (2.00 L) under a dry argon atmosphere was added triethylamine (1.99 g, 19.7 mmol). The solution was cooled to -45° C and diphenylphosphorylazide (6.49 g, 23.8 mmole) was then added dropwise over 5 min.

- 5 The solution was stirred for 22 h while allowing to warm slowly to room temperature. The solution was then stored at -35 °C for an additional 44 h. The solvent was removed *in vacuo* and residual DMF was removed by co-evaporation with H₂O (2 x 300 ml). The residue was dissolved in ethyl acetate (1.5 L), was washed with 0.01 N HCl (2 x 500 ml), saturated NaHCO₃ (2 x 500 ml) and saturated
- 10 NaCl (500 ml), and was dried (MgSO₄). The solvent was removed *in vacuo* to give the crude product as a yellow solid which was purified by flash chromatography (silica gel, CHCl₃:MeOH 98:2 (v/v)) to give 3.40 g (50.7 %) of the macrocycle as a 3:5 mixture of 2 conformations: ¹H NMR (DMSO-d₆, 400 MHz) δ 0.64 (m, 1.1H), 0.79 (d, J = 6.5 Hz, 1.91H), 1.50 - 1.88 (m, 19H), 2.41 (s, 3H), 3.26 (m, 1H),
- 15 3.44 (d, J = 15.6 Hz, 1H), 3.61 - 3.82 (m, 2.2H), 3.87 (quint, J = 6.7 Hz, 0.6H), 3.96 - 4.07 (m, 2.2 H), 4.20 (quint, J = 6.7 Hz, 1H), 5.00 (d, J = 12.9 Hz, 0.35H), 5.06 (d, J = 12.6 Hz, 0.65H), 5.22 (d, J = 13.2 Hz, 0.35H), 5.24 (d, J = 12.6 Hz, 0.65H), 6.90 (d, J = 6.4Hz, 0.6H), 7.09 (d, J = 5.9 Hz, 0.4H), 7.28 - 7.45 (m, 7.4H), 7.53(d, J = 6.2 Hz, 0.6H), 7.94 (m, 2H), 8.55 (m, 1H); ¹³C NMR (DMSO-
- 20 d₆, 100 MHz) δ 16.05, 17.88, 18.63, 18.93, 20.89, 23.75, 23.90, 24.30, 24.37, 24.68, 24.72, 27.37, 30.21, 30.27, 31.27, 31.37, 31.50, 32.36, 46.07, 46.11, 50.56, 50.66, 51.47, 51.88, 52.56, 52.80, 57.04, 57.15, 57.75, 57.86, 58.10, 65.82, 66.02, 79.08, 127.17, 127.36, 127.49, 127.58, 127.61, 127.89, 128.06, 128.08, 128.09, 129.66, 136.08, 136.13, 136.91, 137.20, 143.58, 143.61,
- 25 154.79, 155.64, 169.59, 170.02, 170.95, 171.15, 171.64; FAB mass spectrum (NBA+Li) m/z 688 [M + Li]⁺.

3H. Synthesis of 2R,3R,11R,12R-Bis(cyclohexano)-5R,8R-dimethyl-6,9,14-trioxo-1,4,7,10,13-pentaaza-1-p-toluenesulfonyl-cyclopentadecane

- 30 To a stirred solution of 4-benzyloxycarbonyl-2R,3R,11R,12R-bis(cyclohexano)-5R,8R-dimethyl-6,9,14-trioxo-1,4,7,10,13-pentaaza-1-p-toluenesulfonyl-cyclopentadecane prepared as in example 3G (3.30 g, 4.84 mmole) in a mixture of methanol (60 ml) and H₂O (30 ml) under an argon atmosphere was added ammonium formate (968 mg) and palladium black (484 mg) and the resulting
- 35 mixture was refluxed for 1 h and then stirred for 21 h at room temperature. The

-40-

catalyst was removed by filtration and the solvent was removed *in vacuo* to give 2.69 g of a white solid. Residual ammonium formate was decomposed by addition of palladium black (1 g) to a solution of the crude product in methanol (50 ml) and refluxing for 17 h. Following filtration and removal of the solvent *in vacuo*, the product was purified by flash chromatography (silica gel, CHCl₃:MeOH:aq. NH₄OH, 97.5:2.5:1 (v/v) to give 2.50 g (94.3% yield) of the macrocycle as a colorless solid: ¹H NMR (C₆D₆, 70 °C, 400 MHz) δ 0.65 - 1.40 (m, 18 H), 1.60 (m, 1H), 1.69 (d, J = 7.0 Hz, 3H), 1.90 (s, 3H), 2.31 (br m, 1H), 2.85 (br s, 1H), 3.23 - 3.46 (m, 3H), 3.58 - 3.81 (m, 3H), 4.21 (m, 1H), 6.92 (d, J = 7.9 Hz, 2H), 7.03 (br s, 1H), 7.32 (br d, J = 6.0 Hz, 1H), 7.54 (br s, 1H), 7.67 (d, J = 8.1 Hz, 2H); ¹³C NMR (C₆D₆, 70 °C, 100 MHz) δ 17.79, 18.25, 20.68, 24.30, 24.46, 24.82, 25.36, 28.71, 32.06, 33.15, 49.25, 52.38, 53.17, 54.26, 54.43, 55.47, 64.60, 127.34, 129.63, 138.26, 143.43, 169.48, 172.81, 175.79; FAB mass spectrum (NBA+Li) m/z 554 [M + Li]⁺.

15

3l. Synthesis of 2R,3R,11R,12R-Bis(cyclohexano)-5R,8R-dimethyl-1,4,7,10,13-pentaazacyclopentadecane

To a stirred solution of 2R,3R,11R,12R-bis(cyclohexano)-5R,8R-dimethyl-6,9,14-trioxo-1,4,7,10,13-pentaaza-1-p-toluenesulfonyl-cyclopentadecane prepared as in Example 3H (2.45 g, 4.47 mmole) in anhydrous THF (90 ml) was added a solution of 1.0 M LiAlH₄ (45.0 ml, 45.0 mmole) dropwise over 5 minutes and the bright yellow solution was refluxed for 15 h (by which time it had become heterogeneous). The mixture was then diluted by the addition of THF (200 ml) and was quenched by the dropwise addition of a 10% Na₂SO₄ solution (10 ml) while cooling in an ice bath. The mixture was then filtered and the solids were stirred with anhydrous THF (300 ml) and filtered again. The solids were then refluxed with anhydrous THF (300 ml) for 2 h and stirred overnight at room temperature. Following filtration, the solvent was removed from the extracts which were combined and extracted with ethyl ether. The solvent was removed from the yellow filtrate and the residue was dried by azeotrope with toluene (50 ml) and then hexanes (50 ml) to give 1.67 g of a viscous yellow oil. The product was dissolved in hexanes, leaving behind a small amount of oily byproduct. The solution was concentrated *in vacuo* to give 1.59 g of a yellow oil. The crude product was purified by crystallization from acetonitrile to give 324 mg (20.6% yield) of colorless needles: mp (under argon) 98 - 100 °C; ¹H NMR (C₆D₆, 400 MHz) δ 0.68 (m, 2H), 0.94 (d, J = 6.2 Hz, 3H), 0.97

-41-

- 1.11 (m, 8H), 1.12 (d, $J = 6.4$ Hz, 3H), 1.56 (m, 4H), 1.90 - 2.15 (m, 10H), 2.34 - 2.50 (m, 6H), 2.66 (m, 1H), 2.76 br d, $J = 9.9$ Hz, 1H), 2.94 (br m, 1H), 3.01 (br s, 2H); ^{13}C NMR (C_6D_6 , 100 MHz) δ 17.31, 19.10, 25.33, 25.43, 25.64, 25.80, 31.73, 31.96, 32.30, 32.39, 46.74, 47.79, 52.19, 53.61, 54.71, 57.30, 61.10, 62.38, 62.99; FAB mass spectrum (NBA - HCl) m/z 352 $[\text{M} + \text{H}]^+$.

3I. Synthesis of [Manganese(II)dichloro(2R,3R,11R,12R-bis(cyclohexano)-5R,8R-dimethyl-1,4,7,10,13-pentaazacyclopentadecane)]

- To a stirred solution of MnCl_2 (72 mg, 0.57 mmole) in anhydrous methanol was added 2R,3R,11R,12R-bis(cyclohexano)-5R,8R-dimethyl-1,4,7,10,13-pentaazacyclopentadecane prepared as in Example 3I (201 mg, 0.572 mmole) under a dry nitrogen atmosphere and the solution was refluxed for 1 h. The solvent was removed *in vacuo* and THF (30 ml) was added. To the suspension was added ethanol (1 ml), causing the solid to dissolve. The solution was filtered through CeliteTM and concentrated to a volume of 5 ml. Ethyl ether (40 ml) was then added to the hot solution and the solution was allowed to cool to room temperature while standing for 3 days. The colorless needles were filtered to give 172 mg (35.9% yield) of the product: FAB mass spectrum (NBA) m/z (relative intensity) 443/441 (100/32) $[\text{M} - \text{Cl}]^+$; Anal. Calcd. for $\text{C}_{20}\text{H}_{43}\text{N}_5\text{MnCl}_2$: C, 50.10; H, 9.04; N, 14.61. Found: C, 50.31; H, 8.61; N, 14.66.

Example 4

- 4A. Preparation of 2,3-trans-cycloheptano-5,15-dioxo-7,10,13-tris(p-toluenesulfonyl)-1,4,7,10,13-pentaazacyclopentadecane.

- Two 5 L round bottomed flasks were each charged with 1,2-diaminocycloheptane dihydrochloride (3.49 g, 17.35 mmol), prepared according to (1) from cycloheptanedioxime (2), triethylamine (9.67 ml, 69.4 mmol), and 3,6,9-tris-(p-toluenesulfonyl)-3,6,9-triazaundecanedioic acid (11.83 g, 17.35 mmol) in 3.5 L of degassed DMF. A dry ice-acetone bath was placed around the flask and the solution was cooled to -40°C . Diphenylphosphoryl azide (8.97 ml, 41.6 mmol) was added to the stirred solution over 10 min. The reaction was allowed to slowly warm to room temperature over 16 h. The reaction mixtures were then concentrated together, and the residue was taken up in methylene chloride and washed several times with water. Crystals formed in the methylene chloride layer and were collected to yield 4.27 g of *trans* isomer. The filtrate was washed with aqueous NaHCO_3 .

-42-

aqueous KHSO_4 , water, brine, dried over MgSO_4 , filtered, and concentrated slightly.

A second and third crop of crystals were collected from methylene chloride yielding 8.94 g and 4.48 g of *trans* isomer. The remaining filtrate was concentrated and purified by column chromatography using 1% MeOH in CH_2Cl_2 as eluent. Fractions

- 5 were collected to yield 1.0 g of pure *cis* isomer and 2.7 g of both *cis* and *trans* isomers. A total of 21.4 g of both isomers was collected (79.5 %) including 17.7 g of pure *trans* isomer (66%). 2,3-*trans*-cycloheptano-5,15-dioxo-7,10,13-tris-(p-toluenesulfonyl)-1,4,7,10,13-pentaazacyclopentadecane: ^1H NMR (400 MHz, DMSO- d_6): δ 1.64 - 1.30 (m, 10H), 2.39 (s, 6H), 2.40 (s, 3H), 3.15 - 2.90 (m, 8H), 3.48 (d, J = 16.1 Hz, 2H), 3.75 - 3.65 (m, 2H), 3.97 (d, J = 16.1 Hz, 2H), 7.35 - 7.31 (m, 6H), 7.50 (d, J = 8.1 Hz, 2H), 7.70 (d, J = 8.1 Hz, 4H), 7.92 (d, J = 7.8 Hz, 2H). ^{13}C NMR (100 MHz, DMSO): δ 20.88, 23.96, 27.99, 31.94, 46.29, 46.51, 50.45, 54.03, 126.67, 127.16, 129.55, 129.90, 134.82, 136.13, 143.21, 143.61, 166.01.
- 10
- 15

References

- (1) Belcher, R.; Hoyle, W.; West, T. S. *J. Chem. Soc.* 1961, 667-670.
(2) Vander Haar, R. W.; Voter, R. C.; Banks, C. V. *J. Org. Chem.* 1949, 14, 836-838.

20

4B. Preparation of 2,3-*trans*-cycloheptano-1,4,7,10,13-pentaazacyclopentadecane

Following the procedure used in Example 2D, 2,3-*trans*-cycloheptano-5,15-dioxo-7,10,13-tritosyl-1,4,7,10,13-pentaazacyclopentadecane (6.00 g, 7.75 mmol) was reduced with LiAlH_4 in DME to yield 2,3-*trans*-cycloheptano-1,4,7,10,13-

- 25 pentaazacyclopentadecane. Recrystallization from CH_3CN provided 808 mg of fine needle crystals (37%). ^1H NMR (400 MHz, CDCl_3): δ 1.31 - 1.46 (m, 4H), 1.48 - 1.56 (m, 2H), 1.60 - 1.71 (m, 2H), 1.74 - 1.84 (m, 2H), 2.01 (br s, 5H), 2.18 - 2.25 (m, 2H), 2.45 - 2.54 (m, 2H), 2.57 - 2.77 (m, 10H), 2.79 - 2.86 (m, 2H), 2.91 - 2.98 (m, 2H). ^{13}C NMR (100 MHz, CDCl_3): δ 64.98, 48.73, 48.37, 47.98, 46.86, 29.82, 28.50, 24.02. Anal. Calcd for $\text{C}_{15}\text{H}_{33}\text{N}_5$: C, 63.56; H, 11.73; N, 24.71. Found: C, 63.46; H, 11.66; N, 24.72.
- 30

35

Example 55A. N-p-Toluenesulfonyliminodiacetic acid

- 5 Iminodiacetic acid (200 g, 1.50 mol) was dissolved in water (3 L) and sodium hydroxide (180 g, 4.5 mol) was added in small portions with stirring. After stirring for 15 minutes, p-toluenesulfonyl chloride (286.5 g, 1.5 mol) was added in small portions. The reaction mixture was heated to 80 °C for one hour, giving a clear solution. At the end of this time, the reaction solution was cooled in an ice bath to 10 - 15 °C, and concentrated HCl (110 ml) was added, precipitating the product as a
- 10 white solid. This solid was filtered and washed with cold water, then dried in a vacuum oven at 55 °C, giving 207.7 g (0.723 mol, 48.2% yield): mp 195 - 196 °C (lit.* mp 191 °C); ¹H NMR (DMSO-d₆, 300 MHz) δ 2.38 (s, 3H), 3.66 (s, 4H), 7.39 (d, J = 8.7 Hz, 2H), 7.65 (d, J = 8.1 Hz); ¹³C NMR (DMSO-d₆, 75 MHz) δ 20.90, 53.98, 126.55, 129.79, 136.70, 143.17, 171.77.

15

*J.M. Lehn, J. Simon, J. Wagner, *Nouv. J. Chim.* (1977) 1, 77-84.

5B. Synthesis of Dimethyl 2S,10S-dimethyl-6-p-toluenesulfonyl-4,8-dioxo-3,6,9-triazaundecanedioate

- 20 N-Tosyliminodiacetic acid (28.76 g, 100.1 mmol) was dissolved in N,N-dimethylacetamide (1 L). The solution was degassed and placed under argon. Hydroxybenzotriazole hydrate (41.30 g, 269.8 mmol), L-alanine methyl ester hydrochloride (35.20 g, 252.1 mmol) and triethyl amine (35.4 ml, 254 mmol) were added, then 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (51.8 g,
- 25 270.2 mmol). The heterogeneous reaction was stirred overnight. The solvent was then removed under reduced pressure. Ethyl acetate (500 ml) was added, and water (250 ml) was added, with swirling, to dissolve the residue. The organic phase was washed with saturated NaHCO₃ (2 x 250 ml), 1 M HCl (2 x 250 ml), then saturated NaCl (250 ml). After drying (Na₂SO₄), the solution was concentrated to a white
- 30 foam, 36.5 g, 79.8 mmol, 79.7% yield; mp 144-146 °C; ¹H NMR (CDCl₃, 400 MHz) δ 1.39 (d, J = 7.3 Hz, 6H), 2.41 (s, 3H), 3.71 (s, 6H), 3.87 (s, 4H), 4.50 (quint, J = 7.4 Hz, 2H), 7.30 (d, J = 8.1 Hz, 2H), 7.61 (d, J = 7.3 Hz, 2H), 7.40 (d, J = 8.3 Hz, 2H); ¹³C NMR (CDCl₃, 100 MHz) δ 17.60, 21.57, 48.28, 52.43, 52.63, 127.76, 129.75, 134.63, 144.29, 167.80, 173.04. HRMS. Calcd for
- 35 C₁₉H₂₇N₃O₈S: 457.1519. Found: 457.1538.

5C. Synthesis of 2S,10S-Dimethyl-6-p-toluenesulfonyl-4,8-dioxo-3,6,9-triazaundecanedioic acid

- Dimethyl 2S,10S-dimethyl-6-p-toluenesulfonyl-4,8-dioxo-3,6,9-triazaundecanedioate (25.0 g, 54.6 mmol) was dissolved in 1:1 THF: methanol (300 ml) and stirred while sodium hydroxide solution (2.5 M, 75 ml, 187.5 mmol) was added. After stirring overnight, tlc (silica, developed in 1:9 methanol: dichloromethane) showed that some starting material was still present ($R_f = 0.46$), so additional sodium hydroxide solution (40 ml, 100 mmol) was added, and the reaction mixture was heated for 20 minutes at 40°C, and stirred one hour longer at room temperature. At the end of this time, water (150 ml) was added, and the resulting mixture was washed with ether (3 x 60 ml). The pH of the aqueous layer was adjusted to about 2 by the addition of 1 M KHSO_4 . The aqueous layer was extracted with ethyl acetate (3 x 100 ml). After drying (MgSO_4), the ethyl acetate extract was stripped off and the residue was recrystallized from ethyl acetate - ether, giving the diacid as a white solid, 17.86 g (41.6 mmol, 76% yield): mp 119-120 °C; ^1H NMR ($\text{DMSO}-d_6$, 400 MHz) δ 1.27 (d, $J = 7.25$ Hz, 6H), 2.40 (s, 3H), 3.91 (AB quartet, $J = 17.3$ Hz, $\Delta\nu = 21.1$ Hz, 4H), 4.19 (quintet, $J = 7.1$ Hz, 2H), 7.40 (d, $J = 8.1$ Hz, 2H), 7.71 (d, $J = 7.8$ Hz, 2H), 8.70 (d, $J = 7.25$ Hz, 2H); ^{13}C NMR ($\text{DMSO}-d_6$, 75 MHz) δ 17.11, 20.91, 47.49, 51.20, 127.17, 129.55, 135.43, 143.37, 167.91, 173.54; HRMS. Calcd for $\text{C}_{17}\text{H}_{23}\text{N}_3\text{O}_8\text{S}$: 429.1206. Found: 429.1195. Elemental analysis. Calcd: C, 47.55%; H, 5.40%; N, 9.78%. Found: C, 47.42%; H, 5.41%; N, 9.75%.

5D. Synthesis of 2S,9S-Dimethyl-5R,6R-cyclohexano-3,8,11,15-tetraoxo-13-p-toluenesulfonyl-1,4,7,10,13-pentaazacyclopentadecane

- To a solution of 2S,10S-dimethyl-6-p-toluenesulfonyl-4,8-dioxo-3,6,9-triazaundecanedioic acid (9.66 g, 22.49 mmol) in degassed DMF (4.5 L), triethyl amine (6.2 ml, 45 mmol) and 1R,2R-diaminocyclohexane (2.57 g, 22.5 mmol) were added. The stirred reaction mixture was cooled to -20 °C, and diphenylphosphoryl azide (11.6 ml, 53.8 mmol) was added. After stirring overnight, during which time the reaction reached room temperature, the solvent was removed under reduced pressure. The residue was dissolved in ethyl acetate, and washed with sodium bicarbonate (1:1 saturated solution: water), potassium hydrogen sulfate (5%), and saturated sodium chloride solution. After drying over MgSO_4 and filtration, the solvent was removed under reduced pressure, and the residue was recrystallized from

-45-

acetonitrile as a white crystalline solid, 9.41 g, 18.54 mmol, 82.4% yield. mp 360 °C (dec); ¹H NMR (DMSO-d₆, 400 MHz) δ 1.16, 1.21 (d + m, J = 7.0 Hz, 10H), 1.63 (br s, 2H), 1.90 (m, 2H), 2.42 (s, 3H), 3.41 (br s, 2H), 3.65 (d, J = 16.4 Hz, 2H), 3.91 (d, J = 16.4 Hz, 2H), 4.22 (quint, J = 7.3 Hz, 2H), 7.47 (d, J = 8.1 Hz, 2H), 7.73, 7.77 (d + br s, J = 8.1 Hz, 4H), 8.57 (d, J = 7.8 Hz, 2H); ¹³C NMR (DMSO-d₆, 100 MHz) δ 16.93, 20.91, 24.26, 31.24, 48.44, 52.79, 52.88, 54.84, 127.11, 129.95, 134.33, 143.87, 168.46, 171.59. HRMS. Calcd for C₂₃H₃₃N₅O₆S: 507.2151. Found: 507.2145. Elemental analysis. Calcd: C, 54.42%; H, 6.55%; N, 13.80%. Found: C, 54.15%; H, 6.55%; N, 13.56%.

10

5E. Synthesis of 2S,9S-Dimethyl-5R,6R-cyclohexano-1,4,7,10,13-

pentaazacyclopentadecane

2S,9S-Dimethyl-5R,6R-cyclohexano-3,8,11,15-tetraoxo-13-p-toluenesulfonyl-1,4,7,10,13-pentaazacyclopentadecane (7.0 g, 13.8 mmol) was suspended in 1,2-dimethoxyethane (DME, 350 ml) under argon, and lithium aluminum hydride (0.5 M in DME, 275 ml, 138 mmol) was added rapidly via cannula. The resulting mixture was refluxed overnight, then allowed to cool to room temperature. The reaction mixture was then cooled to -40 °C and water (5.2 ml) was very cautiously and slowly added over a 12 minute period, followed by 15% NaOH (5.2 ml), and water (15.7 ml). THF (400 ml) was added after about 20 minutes and stirring was continued. The cold bath was removed fifteen minutes later. After stirring a few minutes longer, stirring was stopped and the mixture was allowed to settle. The supernatant was decanted via cannula into an argon filled flask, and was subsequently filtered using a cannula tipped (paper) filter. Additional THF was added to the white residue, and stirring was started. After a few hours, the supernatant was removed as before, and filtered. This process was repeated once more using 1 L THF. The combined THF filtrates were stripped under reduced pressure. The residue was extracted with hot hexane, and filtered under argon, and the stripped residue was recrystallized from dry acetonitrile, 2.74 g, 9.21 mmol, 66.8% yield, as a white crystalline solid, mp (under N₂) 179-180 °C; ¹H NMR (C₆D₆, 400 MHz) δ 0.85 (m, 2H), 0.96 (d, J = 6.2 Hz, 6H), 1.11 (m, 2H), 1.40 - 1.66 (m, 4H), 1.71 - 2.08 (m, 9H), 2.42 (m, 2H), 2.55 - 2.66 (m, 6H), 2.72 - 2.83 (m, 4H); ¹³C NMR (C₆D₆, 100 MHz) δ 19.34, 25.49, 32.41, 46.92, 49.03, 54.09, 55.28, 63.82; HRMS. Calcd for C₁₆H₃₅N₅: 297.2892. Found: 297.2889. Elemental analysis. Calcd: C, 64.60%; H, 11.86%; N, 23.54%. Found: C, 64.55%; H, 11.77%; N, 23.54%.

Example 66A. Synthesis of *cis*-5,6-Cyclohexano-3,8-dioxo-1,10,13-tris(p-toluenesulfonyl)-1,4,7,10,13-pentaazacyclopentadecane

- 5 A stirred solution of 3,6,9-tris(p-toluenesulfonyl)-3,6,9-triazaundecanedioic acid (15.3 g, 22.5 mmole), *cis*-1,2-diaminocyclohexane (2.57 g, 22.5 mmole) and triethylamine (4.55 g, 45.0 mmole) in degassed anhydrous DMF (4.50 L) under a dry argon atmosphere was cooled to -40 °C and diphenylphosphorylazide (14.9 g, 54.0
- 10 mmole) was added dropwise over 10 minutes. The mixture was allowed to warm to room temperature while stirring for 15 h and the solvent was removed *in vacuo*. The residue was dissolved in ethyl acetate (2.0 L), washed with 0.1 N HCl (2 x 500 ml), saturated NaHCO₃ solution (2 x 500 ml) and saturated NaCl solution (500 ml), and was dried (MgSO₄). After filtration, the solution was concentrated to a volume of
- 15 250 ml and ethyl ether was added periodically as the crystallization progressed. The crystals were filtered, washed with ether and dried to give 15.0 g of crude product. Recrystallization from CH₂Cl₂ - methanol and flash chromatographic purification of the filtrate (silica gel, CHCl₃: MeOH, 97.5:2.5 (v/v)) gave a total of 11.54 g (67.5% yield) of the macrocycle: mp 256 - 257 °C; ¹H NMR (CDCl₃, 400 MHz) δ 1.53 (br
- 20 s, 6H), 1.78 (br s, 2H), 2.45 (s, 9H), 3.18 (m, 2H), 3.37 (m, 2H), 3.46 (d, J = 16.7 Hz, 2H), 3.51 (m, 4H), 3.82 (d, J = 16.9 Hz, 2H), 4.16 (br s, 2H), 6.84 (d, J = 7.3 Hz, 2 H), 7.36 (d, J = 8.3 Hz, 6H), 7.71 (d, J = 8.3 Hz, 4H), 7.75 (d, J = 8.3 Hz, 2H); ¹³C NMR (CDCl₃, 100 MHz) δ 21.61, 22.04, 28.14, 49.46, 49.58, 51.52, 54.60, 127.61, 127.75, 130.00, 130.11, 133.64, 135.29, 144.07, 144.60,
- 25 168.20; FAB mass spectrum (NBA - Li) *m/z* 766 [M + Li]⁺.

6B. Synthesis of *cis*-2,3-Cyclohexano-1,4,7,10,13-pentaazacyclopentadecane

- To a stirred suspension of *cis*-5,6-cyclohexano-1,10,13-tris(p-toluenesulfonyl)-1,4,7,10,13-pentaazacyclopentadecane-3,8-dione prepared as in
- 30 example 6A (13.5 g, 17.8 mmole) in anhydrous THF (340 ml) under a dry argon atmosphere was added a solution of 1.0 M LiAlH₄ in THF (222 ml, 222 mmole) dropwise over 30 minutes. The yellow homogeneous solution was refluxed for 17 h, by which time it had become heterogeneous. The mixture was then cooled to room temperature in a water bath and H₂O (7.1 ml) was added dropwise over 5 minutes,
- 35 followed 5 minutes later by a 15% solution of NaOH (7.1 ml) and then 5 minutes

-47-

- later by H₂O (20.7 ml). The slurry was filtered and the solids were refluxed with THF (2 x 500 ml) for 30 minutes and filtered. Then the solids were stirred with a mixture of THF (250 ml) and MeOH (250 ml) at room temperature for 2 days and then refluxed for 3 h. The mixture was filtered and the residue was redissolved in THF and filtered again. The filtrates were combined and the solvent was removed *in vacuo*. The crude product was crystallized from acetonitrile to give 2.22 g (46.3%) of the product which contains 5% of the *trans* isomer. Recrystallization from hexanes and then acetonitrile reduced the amount of *trans* isomer to 2%, giving 1.92 g (32.4% yield) of the product as colorless needles: mp (under argon) 115 - 116° C; ¹H NMR (C₆D₆, 400 MHz) δ 1.24 (m, 4H), 1.43 (br s, 5H), 1.70 (m, 4H), 2.43 (m, 2H), 2.49 - 2.62 (m, 14 H), 2.68 (m, 2H); ¹³C NMR (CDCl₃, 100 MHz) δ 22.21, 27.75, 46.49, 48.15, 48.58, 48.97, 57.24; CI mass spectrum (CH₄) m/z 270 [M + H]⁺.

15

Example 7

- 7A. 1,7-Bis(p-toluenesulfonyl)-2R,3R-cyclohexano-5-oxo-1,4,7-triazaheptane
N-p-Toluenesulfonyl-1R,2R-diaminocyclohexane (15 g, 55.9 mmol) was dissolved in DMF (550 ml), and triethylamine (9.4 ml, 61.1 mmol) and N-p-toluenesulfonyl glycine (15.4 g, 67.1 mmol) were added. The solution was cooled to -20 °C, and diphenylphosphoryl azide (14.5 ml, 67.2 mmol) was added dropwise over 8 minutes. The reaction was allowed to warm to room temperature overnight. After removal of solvent under reduced pressure, the residue was dissolved in dichloromethane (200 ml) and washed with water (50 ml), sodium bicarbonate (1:1 saturated: water, 50 ml), 5% KHSO₄ (50 ml), then saturated NaCl (50 ml). The organic layer was dried (MgSO₄), filtered, and the solvent was removed under reduced pressure, and the residue was recrystallized from dichloromethane-hexanes, 21.76 g, (45.37 mmol), 81% yield, as a white crystalline solid, mp 96-97 °C; ¹H NMR (CDCl₃, 400 MHz) δ 1.0 - 1.27 (m, 4H), 1.52 - 1.71 (m, 3H), 1.81 - 1.92 (br s, 1H), 2.38, 2.39 (2 s, 6H), 2.96 (m, 1H), 3.4 - 3.68 (m, 3H), 5.68 (d, J = 7.8 Hz, 1H), 5.98 (t, J = 6.3 Hz, 1H), 6.76 (d, J = 8.9 Hz), 7.26 (d, J = 7.8 Hz, 4 H), 7.72 (m, 4H); ¹³C NMR (CDCl₃, 100 MHz) δ 21.51, 24.50, 24.74, 31.92, 33.21, 46.09, 52.98, 57.95, 126.88, 127.24, 129.69, 129.86, 136.19, 138.55, 143.24, 143.78, 169.26; HRMS. Calcd for C₂₂H₂₉N₃O₅S₂: 479.1548. Found: 479.1549.

7B. Preparation of Dimethyl 3,9-bis(p-toluenesulfonyl)-4R,5R-cyclohexano-7-oxo-3,6,9-triazaundecandioate

- 5 1,7-Bis(p-toluenesulfonyl)-2R,3R-cyclohexano-5-oxo-1,4,7-triazaheptane (5.0 g, 10.4 mmol) was dissolved in DMF (100 ml) under argon, and methyl bromoacetate was added (5.9 ml, 62.3 mmol). The reaction progress was monitored by tlc (on silica, developed in 10% methanol, 90% dichloromethane, $R_f = 0.57$ for starting material, $R_f = 0.65$ for diester product) and additional reagents were added as necessary. After cooling the solution to 0 °C, sodium hydride (1.06 g, 44.2 mmol)
- 10 was added with stirring. At the end of the addition, the ice bath was removed, and the reaction was stirred for one hour at room temperature. Heating was begun in a water bath, for 2.75 h, at an internal temperature of 35 to 40 °C. Twenty minutes before the end of the heating period, additional methyl bromoacetate (2 ml, 21.1 mmol) was added. The reaction was stirred overnight at room temperature. The
- 15 reaction was placed in an ice bath, and additional sodium hydride (0.26 g, 10.8 mmol) and methyl bromoacetate (1.0 ml, 10.6 mmol) were added, and the ice bath was removed. Three hours later, water (200 ml) was added to the reaction mixture, and the solvents were removed under reduced pressure. The residue was dissolved in ethyl acetate (150 ml), washed with water (3 x 80 ml), saturated sodium chloride
- 20 solution (50 ml), and dried over $MgSO_4$. After filtration, the solvent was removed under reduced pressure. The product was used in the next step without further purification. MS m/z (relative intensity) 630 (100, $[M + Li^+]$).

7C. Preparation of 3,9-Bis(p-toluenesulfonyl)-4R,5R-cyclohexano-7-oxo-3,6,9-triazaundecanedioic acid

- 25 Dimethyl 3,9-bis(p-toluenesulfonyl)-4R,5R-cyclohexano-7-oxo-3,6,9-triazaundecandioate, from the previous preparation, without additional purification, was dissolved in methanol (40 ml) and stirred while aqueous sodium hydroxide (2.5N, 25 ml, 62.5 mmol) was added. After stirring overnight, tlc indicated the
- 30 presence of starting material. Additional sodium hydroxide solution (8 ml of 2.5 M, 20 mmol) was added and the reaction flask was heated in a 50 °C water bath for 4.5 hours. At the end of this time, the solvent was removed under reduced pressure. Water was added and the resulting solution was washed twice with ether (30 ml) and the pH was adjusted to about 3 by the addition of 1M $KHSO_4$. The aqueous layer
- 35 was extracted with ethyl acetate (3 x 50 ml). The combined ethyl acetate layers were dried over $MgSO_4$, filtered and the solvent was removed under reduced pressure.

- The residue was chromatographed twice on silica, eluting with 8% methanol, 1% acetic acid, 91% dichloromethane. The fractions containing product ($R_f = 0.29$, on a silica plate, developed in 1% acetic acid, 10% methanol, 89% dichloromethane) were combined and chromatographed on silica initially eluting with 1% acetic acid, 2% methanol, 97% dichloromethane, and later eluting with 1% acetic acid, 4% methanol, 95% dichloromethane. Product containing fractions were combined, and the solvents removed under reduced pressure. Toluene was added to the residue, and the resulting mixture was concentrated, and the process was repeated. The product was obtained as an off-white solid, 4.58 g, 7.69 mmol, 73.7% (for two steps): mp 265-268 °C
- 5 (dec). ^1H NMR (CD_3OD , 400 MHz) δ 1.00 - 1.38 (m, 4H), 1.46 (m, 1H), 1.60 (m, 2H), 1.96 (m, 1H), 2.40 (s, 3H), 2.42 (s, 3H), 3.33 (m, 1H), 3.68 (m, 1H), 3.75 - 4.2 (m, 7H), 7.33 (d, $J = 8.1$ Hz, 2H), 7.39 (d, $J = 8.1$ Hz, 2H), 7.75 (d, $J = 8.1$ Hz, 2H), 7.78 (d, $J = 8.3$ Hz, 2H). Electrospray mass spectrum m/z (relative intensity) 618 (100, $[\text{M} + \text{Na}]^+$), 596 (25, $[\text{M} + \text{H}]^+$).

15

7D. Preparation of 2R,3R,8R,9R-Bis(cyclohexano)-3,13-bis(p-toluenesulfonyl)-6,11,15-trioxo-1,4,7,10,13-pentaazacyclopentadecane.

- Triethylamine (1.12 ml, 8.05 mmol), 1R,2R-diaminocyclohexane (0.46 g, 4.02 mmol), and 7-oxo-3,9-bis(p-toluenesulfonyl)-4R,5R-cyclohexano-3,6,9-triazaundecanedioic acid (2.4 g, 4.0 mmol) were dissolved in degassed DMF (1000 ml) and cooled to -20 °C under argon. Diphenylphosphoryl azide (2.08 ml, 9.66 mmol) was added, and the reaction was stirred overnight, rising to room temperature. The solvent was removed under reduced pressure. Dichloromethane was added to the residue, precipitating a white powder. The filtrate was stripped, then dissolved in
- 20 dichloromethane (200 ml) and washed successively with single portions of sodium bicarbonate, 5% potassium bisulfate, and saturated sodium chloride solution. After drying (MgSO_4) and filtering, the solvent was removed under reduced pressure.
- Recrystallization from acetonitrile gave successive crops totalling 1.62 g, 2.40 mmol, 60% yield: white crystals; mp 298-299 °C; ^1H NMR (CDCl_3 , 400 MHz) δ 0.57 (m, 2H), 1.11 - 1.48 (m, 6H), 1.55 - 1.89 (m, 5H), 1.98 - 2.52 (s + m, 9H), 3.15 (d, $J = 14.5$ Hz, 1H), 3.41 - 3.76 (m, 6H), 4.14 (m, 1H), 4.26 (d, $J = 18.3$ Hz, 1H), 4.63 (d, $J = 17.2$ Hz, 1H), 5.93 (d, $J = 9.9$ Hz), 6.28 (d, $J = 5.4$ Hz, 1H), 7.27 (m, 4H), 7.53 (d, $J = 8.1$ Hz, 1H), 7.75 (d, $J = 8.1$ Hz, 1H), 8.64 (d, $J = 8.6$ Hz, 1H);
- 30 ^{13}C NMR (CDCl_3 , 100 MHz) δ 21.51, 24.23, 24.34, 24.89, 25.52, 28.48, 31.91, 32.75, 33.68, 46.80, 50.78, 51.43, 53.60, 54.21, 54.44, 62.04, 126.64, 127.81,
- 35

-50-

129.86, 130.11, 135.77, 136.24, 144.13, 144.17, 167.32, 168.17, 168.27; MS m/z (relative intensity) 680 (100 [M + Li]⁺); Elemental analysis. Calcd: C, 57.04%; H, 6.43%; N, 10.39%. Found: C, 56.99%; H, 6.57%; N, 10.27%.

5

Example 88A. Preparation of Dimethyl 2R-methyl-3-azapentandioate.

L-Alanine methyl ester hydrochloride (24.24 g, 173.7 mmol) in 600 ml of DMF was placed in a 2L round bottomed flask under argon atmosphere.

- 10 Triethylamine (48.4 ml, 347 mmol) was added to the solution and precipitate began to form. Methyl bromoacetate (17.0 ml, 173.7 mmol) was added dropwise to the mixture, and the mixture was stirred at room temperature. After 16 h, the mixture was concentrated, EtOAc and water were added and the layers were combined and separated. The EtOAc layer was washed with water and brine, dried over Na₂SO₄,
15 filtered and concentrated to yield 20.0 g of light tan oil (66%). Dimethyl 2S-methyl-3-azapentandioate was prepared from D-alanine methyl ester hydrochloride, in the same manner. ¹H NMR (CDCl₃, 300 MHz) δ 1.30 (d, J = 6.9 Hz, 3H), 2.03 (s, 1H), 3.30 - 3.49 (m, 3H), 3.68 (s, 6H).

20 8B. Preparation of Dimethyl 2S-methyl-3-p-toluenesulfonyl-3-azapentanoate.

Dimethyl 2S-methyl-3-azapentanoate (19.7 g, 112 mmol) was gradually added to a mixture of p-toluenesulfonylchloride (24.0 g, 126 mmol) in 150 ml of pyridine. The resulting light orange solution was placed in an oil bath at 50 °C and stirred for 1 h. The mixture was allowed to cool to room temperature then was cooled
25 in an ice-water bath. Water (250 ml) was added to the mixture and the solution was stirred for a few minutes then concentrated. The residue was taken up in water and EtOAc. The layers were separated and the EtOAc layer was washed with water until the washes were colorless. The EtOAc layer was then washed with brine, dried over MgSO₄, filtered and concentrated to a light tan oil, weight 23.5 g (64%). Dimethyl-

- 30 2R-methyl-3-p-toluenesulfonyl-3-azapentanoate was prepared in the same manner. ¹H NMR (CDCl₃, 300 MHz) δ 1.35 (d, J = 7.2 Hz, 3H), 2.39 (s, 3H), 3.51 (s, 3H), 3.70 (s, 3H), 4.01 (d, J = 19 Hz, 1H), 4.26 (d, J = 19 Hz, 1H), 4.50 (q, J = 7.2, 15 Hz, 1H), 7.27 (d, J = 8.1 Hz, 2H), 7.74 (d, J = 8.4 Hz, 2H); ¹³C NMR (CDCl₃, 100 MHz) δ 16.3, 21.5, 45.3, 52.1, 52.2, 54.5, 127.6, 129.4, 136.4,
35 143.7, 170.2, 171.7; MS (FAB+ DTT:DTE) m/z 330 (M + H), 270 (-CO₂CH₃).

8C. Preparation of 2S-Methyl-3-p-toluenesulfonyl-3-azapentandioic acid

- A solution of dimethyl 2S-methyl-3-p-toluenesulfonyl-3-azapentandioate (23.5 g, 71.3 mmol) in THF (250 ml) was cooled in an ice-water bath. Aqueous NaOH solution (1N, 200 ml) was gradually added to the solution and the mixture was stirred. The ice bath was removed and the mixture was allowed to warm to room temperature. After 1 hour, the mixture was acidified by addition of 250 ml of 1N HCl and concentrated to ~50 ml volume. The aqueous mixture was extracted with 500 ml of EtOAc. The EtOAc layer was washed with 50 ml of water, brine, dried over MgSO₄, filtered and concentrated to give an amber colored oil. CH₂Cl₂ was added to the residue and evaporated to give an off-white solid, 20.6 g. The solid was recrystallized from EtOAc and hexanes to give 19.1 g of spherical crystals (89%). 2R-Methyl-3-p-toluenesulfonyl-3-azapentandioic acid was prepared in the same manner. ¹H NMR (DMSO-d₆, 300 MHz) δ 1.28 (d, J = 7.5 Hz, 3H), 2.36 (s, 3H), 3.96 (d, J = 18 Hz, 1H), 4.08 (d, J = 18 Hz, 1H), 4.26 - 4.38 (m, 1H), 7.35 (d, J = 8.4 Hz, 2H), 7.71 (d, J = 8.4 Hz, 2H), 12.69 (br s, 2H). ¹³C NMR (DMSO-d₆, 100 MHz) δ 16.2, 21.0, 45.8, 54.5, 127.1, 129.5, 136.7, 143.2, 171.1, 172.2.

8D. Synthesis of Dimethyl 2S,5R,10S-trimethyl-4,8-dioxo-6-p-toluenesulfonyl-3,6,9-triazaundecanedioate

- N,N-Dimethylacetamide (450 ml) was degassed, and R-2-methyl-N-p-toluenesulfonyliminodiacetic acid (14.05 g, 46.3 mmol), S-alanine methyl ester hydrochloride (16.47 g, 118.0 mmol), and hydroxybenzotriazole hydrate (19.05 g, 124.4 mmol) were added. A few minutes later, triethylamine (16.5 ml, 118 mmol) and EDC·HCl (24.54 g, 128 mmol) were added and the mixture was stirred overnight. After 15 h, the solvent was removed under reduced pressure, and the residue was briefly placed under vacuum. Ethyl acetate (250 ml) and water (125 ml) were added, and the mixture was agitated briefly. The layers were separated and the aqueous layer was washed with ethyl acetate (100 ml). The combined organic layers were washed successively with saturated sodium bicarbonate (twice), 1 M HCl (twice), and saturated sodium chloride (once). After drying (Na₂SO₄) and filtration, the solvent was removed under reduced pressure, giving a white solid (21.17 g, 44.99 mmol, 97% yield): ¹H NMR (CDCl₃, 400 MHz) δ 1.27 (d, J = 7.0 Hz, 3H), 1.32 (d, J = 7.3 Hz, 3H), 1.40 (d, J = 7.0 Hz, 3H), 2.38 (s, 3H), 3.65 (s, 3H), 3.69 (s, 3H), 3.80 (d, J = 16.9 Hz, 1H), 3.97 (d, J = 16.9 Hz, 1H), 4.25 (q, J =

-52-

7.2 Hz, 1H), 4.33 (quint, $J = 7.3$ Hz, 1H), 4.55 (quint, $J = 7.3$ Hz, 1H), 7.07 (d, $J = 7.5$ Hz, 1H), 7.26 (d, $J = 8.1$ Hz, 2H), 7.78 (d, $J = 8.3$ Hz, 2H), 8.30 (d, $J = 7.3$ Hz, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 14.86, 16.95, 18.00, 21.57, 47.51, 48.32, 48.42, 52.26, 52.42, 56.55, 128.06, 129.77, 135.35, 144.41, 169.12, 170.60, 172.97, 173.09.

8E. Preparation of 2S,5R,10S-Trimethyl-6-p-toluenesulfonyl-4,8-dioxo-3,6,9-triazaundecanedioic acid

Dimethyl 2S,5R,10S-trimethyl-6-p-toluenesulfonyl-4,8-dioxo-3,6,9-triazaundecandioate (21.17 g, 44.90 mmol) was hydrolyzed using the procedure described in Example 8C in 180 ml of THF and using NaOH (3.59 g, 89.79 mmol) in 90 ml of water. A dry foam was obtained, weight 20.9 g. ^1H NMR ($\text{DMSO}-d_6$, 400 MHz) δ 1.06 (d, $J = 7.0$ Hz, 3H), 1.25 (d, $J = 7.2$ Hz, 3H), 1.30 (d, $J = 7.2$ Hz, 3H), 2.40 (s, 3H), 3.75 (d, $J = 17.5$ Hz, 1H), 4.03 (d, $J = 17.5$ Hz, 1H), 3.98 - 4.11 (m, 1H), 4.25 - 4.38 (m, 2H), 7.39 (d, $J = 8.1$ Hz, 2H), 7.77 (d, $J = 8.1$ Hz, 2H), 8.48 (d, $J = 7.5$ Hz, 1H), 8.96 (d, $J = 7.0$ Hz, 1H), 12.56 (br s, 2H); ^{13}C NMR ($\text{DMSO}-d_6$, 100 MHz) δ 15.4, 16.9, 17.6, 21.1, 46.8, 47.7, 47.8, 55.5, 127.5, 129.7, 135.7, 143.7, 169.2, 170.8, 173.7.

8F. Preparation of 3S,6R,11S-Trimethyl-2,5,9,12-tetraoxo-7-p-toluenesulfonyl-1,4,7,10,13-pentaazacyclopentadecane.

Ethylene diamine (151 μl , 2.25 mmol) and triethylamine (629 μl , 4.51 mmol) were added to a solution of 2S,5R,10S-trimethyl-6-p-toluenesulfonyl-4,8-dioxo-3,6,9-triazaundecanedioic acid (1.00 g, 2.25 mmol) in 450 ml of degassed DMF. The solution was cooled to between -40 and -50 $^\circ\text{C}$ under argon atmosphere and diphenylphosphoryl azide (1.16 ml, 5.40 mmol) was added. The reaction was stirred and allowed to slowly warm to room temperature overnight. After ~ 16 h, The reaction mixture was concentrated *in vacuo* and the residue was dissolved in EtOAc and water. The EtOAc layer was separated, washed with water, aq. NaHCO_3 , 0.5 N KHSO_4 , water, brine, and dried over MgSO_4 , filtered, and concentrated to an oil. The oil was dissolved in ~ 15 ml of CH_3CN and filtered. EtOAc and diethyl ether (Et_2O) were added to the concentrated solution and fluffy white solid formed to give 200 mg. The solid was purified by column chromatography on silica gel using 5% MeOH in CH_2Cl_2 as eluent. The product was collected as a white solid, 129 mg (12%). ^1H NMR (CDCl_3 , 400 MHz) δ 1.10 - 1.18 (m, 6H), 1.33 (d, $J = 7.0$ Hz,

-53-

- 3H), 2.38 (s, 3H), 3.17 - 3.35 (m, 3H), 3.38 - 3.49 (m, 1H), 3.88 - 3.96 (m, 1H), 4.17 - 4.30 (m, 3H), 4.32 - 4.42 (m, 1H), 6.86 (br s, 1H), 7.11 (br s, 1H), 7.29 (d, $J = 8.1$ Hz, 2H), 7.69 (br s, 1H), 7.76 (d, $J = 8.1$ Hz, 2H), 8.42 (d, $J = 8.1$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 14.1, 16.4, 17.8, 21.5, 37.9, 38.8, 46.8, 49.7, 50.8, 56.7, 127.8, 130.0, 135.3, 144.7, 170.8, 172.6, 172.9; FAB mass spectrum (NBA+Li) m/z 474 $[\text{M} + \text{Li}]^+$, 480 $[\text{M} - \text{H} + 2\text{Li}]^+$, 516 $[\text{M} + \text{Li} + \text{LiCl}]^+$.

Example 9

- 10 9A. Preparation of 2R,3R-Cyclohexano-6S,9R,14S-trimethyl-5,8,12,15-tetraoxo-10-p-toluenesulfonyl-1,4,7,10,13-pentaazacyclopentadecane.
- A 5 L round bottomed flask was charged with 2S,5R,10S-trimethyl-6-p-toluenesulfonyl-4,8-dioxo-3,6,9-triazaundecanedioic acid (6.65 g, 15.00 mmol), prepared as in Example 8E, in 3 L of anhydrous DMF. The clear solution was cooled in an ice-water bath, and HOBT \cdot H₂O and EDC \cdot HCl were added. After 15 min, 1R,2R-diaminocyclohexane in ~750 ml of DMF was added to the solution via cannula. The addition was complete after ~30 min. The solution was allowed to slowly warm to room temperature overnight. After 16 h, the reaction mixture was concentrated to 20 ml volume and diluted with EtOAc and aqueous NaHCO₃. The layers were separated and the aqueous layer was extracted with EtOAc four times. The EtOAc layers were combined and washed with 0.5 N NaHSO₄, H₂O, and brine, then concentrated. Some precipitate that formed after washing the EtOAc layer with water was rinsed with water, dissolved in MeOH and added to the crude product. The crude product, 7.7 g, was heated in 500 ml of acetonitrile and filtered. The product was collected as a white solid, weight 4.00 g (51%). ^1H NMR (CDCl_3 with 10% CD_3OD , 400 MHz) δ 1.00 - 1.20 (m, 13H), 1.51 - 1.63 (m, 2H), 1.72 - 1.90 (m, 2H), 2.28 (s, 3H), 3.33 - 3.41 (m, 3H), 3.98 - 4.14 (m, 4H), 7.20 (d, $J = 8.0$ Hz, 2H), 7.63 (d, $J = 8.4$ Hz, 2H); ^{13}C NMR (CDCl_3 with 10% CD_3OD , 100 MHz) δ 13.8, 15.6, 17.0, 21.1, 24.3, 24.4, 31.8, 31.9, 45.6, 49.3, 49.6, 52.3, 52.7, 56.8, 127.3, 129.7, 135.3, 144.3, 169.9, 171.3, 171.4, 173.2; FAB mass spectrum (TGL) m/z 522 (M+H).

Example 10Stopped-Flow Kinetic Analysis

- Stopped-flow kinetic analysis has been utilized to determine whether a
- 5 compound can catalyze the dismutation of superoxide (Riley, D.P., Rivers, W.J. and Weiss, R.H., "Stopped-Flow Kinetic Analysis for Monitoring Superoxide Decay in Aqueous Systems," *Anal. Biochem.* 196, 344-349 [1991]). For the attainment of consistent and accurate measurements all reagents were biologically clean and metal-free. To achieve this, all buffers (Calbiochem) were biological grade, metal-free
- 10 buffers and were handled with utensils which had been washed first with 0.1 N HCl, followed by purified water, followed by a rinse in a 10^{-4} M EDTA bath at pH 8, followed by a rinse with purified water and dried at 65°C for several hours. Dry DMSO solutions of potassium superoxide (Aldrich) were prepared under a dry, inert atmosphere of argon in a Vacuum Atmospheres dry glovebox using dried glassware.
- 15 The DMSO solutions were prepared immediately before every stopped-flow experiment. A mortar and pestle were used to grind the yellow solid potassium superoxide (~100 mg). The powder was then ground with a few drops of DMSO and the slurry transferred to a flask containing an additional 25 ml of DMSO. The resultant slurry was stirred for 1/2 h and then filtered. This procedure gave
- 20 reproducibly ~2 mM concentrations of superoxide in DMSO. These solutions were transferred to a glovebag under nitrogen in sealed vials prior to loading the syringe under nitrogen. It should be noted that the DMSO/superoxide solutions are extremely sensitive to water, heat, air, and extraneous metals. A fresh, pure solution has a very slight yellowish tint.
- 25 Water for buffer solutions was delivered from an in-house deionized water system to a Barnstead Nanopure Ultrapure Series 550 water system and then double distilled, first from alkaline potassium permanganate and then from a dilute EDTA solution. For example, a solution containing 1.0 g of potassium permanganate, 2 liters of water and additional sodium hydroxide necessary to bring the pH to 9.0 were
- 30 added to a 2-liter flask fitted with a solvent distillation head. This distillation will oxidize any trace of organic compounds in the water. The final distillation was carried out under nitrogen in a 2.5-liter flask containing 1500 ml of water from the first still and 1.0×10^{-6} M EDTA. This step will remove remaining trace metals from the ultrapure water. To prevent EDTA mist from volatilizing over the reflux arm to
- 35 the still head, the 40-cm vertical arm was packed with glass beads and wrapped with

-55-

insulation. This system produces deoxygenated water that can be measured to have a conductivity of less than 2.0 nanomhos/cm².

- The stopped-flow spectrometer system was designed and manufactured by Kinetic Instruments Inc. (Ann Arbor, MI) and was interfaced to a MAC IICX personal computer. The software for the stopped-flow analysis was provided by Kinetics Instrument Inc. and was written in QuickBasic with MacAdios drivers. Typical injector volumes (0.10 ml of buffer and 0.006 ml of DMSO) were calibrated so that a large excess of water over the DMSO solution were mixed together. The actual ratio was approximately 19/1 so that the initial concentration of superoxide in the aqueous solution was in the range 60-120 μ M. Since the published extinction coefficient of superoxide in H₂O at 245 nm is $-2250 \text{ M}^{-1} \text{ cm}^{-1}$ (1), an initial absorbance value of approximately 0.3-0.5 would be expected for a 2-cm path length cell, and this was observed experimentally. Aqueous solutions to be mixed with the DMSO solution of superoxide were prepared using 80 mM concentrations of the Hepes buffer, pH 8.1 (free acid + Na form). One of the reservoir syringes was filled with 5 ml of the DMSO solution while the other was filled with 5 ml of the aqueous buffer solution. The entire injection block, mixer, and spectrometer cell were immersed in a thermostatted circulating water bath with a temperature of $21.0 \pm 0.5^\circ\text{C}$.
- Prior to initiating data collection for a superoxide decay, a baseline average was obtained by injecting several shots of the buffer and DMSO solutions into the mixing chamber. These shots were averaged and stored as the baseline. The first shots to be collected during a series of runs were with aqueous solutions that did not contain catalyst. This assures that each series of trials were free of contamination capable of generating first-order superoxide decay profiles. If the decays observed for several shots of the buffer solution were second-order, solutions of manganese(II) complexes could be utilized. In general, the potential SOD catalyst was screened over a wide range of concentrations. Since the initial concentration of superoxide upon mixing the DMSO with the aqueous buffer was $\sim 1.2 \times 10^{-4} \text{ M}$, we wanted to use a manganese (II) complex concentration that was at least 20 times less than the substrate superoxide. Consequently, we generally screened compounds for SOD activity using concentrations ranging from 5×10^{-7} to $8 \times 10^{-6} \text{ M}$. Data acquired from the experiment was imported into a suitable math program (e.g., Cricket Graph) so that standard kinetic data analyses could be performed.
- The catalytic rate constant for dismutation of superoxide by the manganese(II) complex of Example 1 was determined from the linear plot of observed rate constants

-56-

(k_{obs}) versus the concentration of the manganese(II) complexes. k_{obs} values were obtained from the liner plots of \ln absorbance at 245 nm versus time for the dismutation of superoxide by the manganese(II) complex. The k_{cat} ($M^{-1}sec^{-1}$) of the manganese (II) complex of Examples 1-3 at pH = 8.1 and 21°C were determined and the results can be found in Table I.

The manganese(II) complexes of fifteen-membered pentaazamacrocyclic ligand in Examples 1-3 are effective catalysts for the dismutation of superoxide, as can be seen from the k_{cat} data in Table I.

10

TABLE I


Example No.	k_{cat} ($M^{-1}sec^{-1}$)
1	3.92×10^7
2	1.29×10^7
3	1.36×10^7

15


-57-

WHAT IS CLAIMED IS:

1. A process for preparing a substituted polyazamacrocyclic comprising contacting (a) a diamine represented by the formula:




- and (b) a dicarboxylic acid or ester thereof represented by the formula:



- or an anhydride thereof; in the presence of a suitable base and a suitable solvent under reactions conditions sufficient to produce said substituted polyazamacrocyclic represented by the formula:

15

-58-



- provided that when an ester of said dicarboxylic acid is used, said suitable base is optional, and when said dicarboxylic acid or an anhydride of said dicarboxylic acid is used, the reaction mixture further comprises a suitable coupling agent; wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₁₁ and R₁₂ are independently selected from hydrogen or alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, alkylcycloalkyl, alkenylcycloalkyl, alkylcycloalkenyl, alkenylcycloalkenyl, heterocyclic, aryl and aralkyl radicals or radicals attached to the α-carbon of α-amino acids; or R₃ or R₄ and R₅ or R₆, and R₇ or R₈ and R₉ or R₁₀ together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms; or R₁ or R₂ and R₃ or R₄, R₅ or R₆ and R₇ or R₈, and R₉ or R₁₀ and R₁₁ or R₁₂ together with the carbon atoms to which they are attached independently form a nitrogen containing heterocycle having 2 to 20 carbon atoms provided that when the nitrogen containing heterocycle is an aromatic heterocycle which does not contain a substituent attached to the nitrogen, the hydrogen attached to the nitrogen in said formula, which nitrogen is also in the macrocycle and the R groups attached to the same carbon atoms of the macrocycle are absent; R₁ and R₂, R₃ and R₄, R₅ and R₆, R₇ and R₈, R₉ and R₁₀, and R₁₁ and R₁₂ together with the carbon atom to which they are attached independently form a saturated, partially saturated or unsaturated ring structure having 3 to 20 carbon atoms, or R₁ and R₂, R₃ and R₄, R₅ and R₆, R₇ and R₈, R₉ and R₁₀, and R₁₁ and R₁₂ independently are =O or =S; and combinations thereof; and wherein R₁₃, R₁₄, R₁₅, R₁₆, R₁₉ and R₂₀ are independently selected from hydrogen or alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cycloalkylalkyl,

-59-

- cycloalkylcycloalkyl, cycloalkenylalkyl, alkylcycloalkyl, alkenylcycloalkyl, alkylcycloalkenyl, alkenylcycloalkenyl, heterocyclic, aryl and aralkyl radicals; or R_{13} and R_{14} , R_{15} and R_{16} , and R_{19} and R_{20} together with the carbon atom to which they are attached independently form a saturated, partially saturated or unsaturated ring structure having 3 to 20 carbon atoms; or
- 5 R_{19} and R_{20} are independently $-OR_{23}$, $-OH$, $-SR_{23}$, $-NR_{23}R_{24}$, $-P(O)(OR_{25})(OR_{26})$ or $-P(O)(R_{25})(OR_{26})$; or
- R_{19} and R_{20} are $=O$, $=S$, $=NR_{23}$, $=N-OH$, $=N-OR_{23}$, $=N-O-C(O)-R_{23}$, or $=CR_{23}R_{24}$; or
- 10 R_{13} or R_{14} and R_{15} or R_{16} together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms, provided that if said diamine has three carbons between the nitrogen atoms, the saturated, partially saturated or unsaturated cyclic has 4 to 20 carbon atoms; or
- 15 R_{13} or R_{14} and R_{19} or R_{20} , or R_{15} or R_{16} and R_{19} or R_{20} together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms; and combinations thereof; wherein R_{17} and R_{18} are independently selected from hydrogen and alkyl or aryl groups;
- 20 wherein R is hydrogen or alkyl or aryl groups; wherein R_{23} and R_{24} are independently selected from alkyl, aralkyl or aryl groups; R_{25} and R_{26} are independently selected from hydrogen, alkyl, aralkyl or aryl groups and
- wherein A , B and C are independently selected from hydrogen, alkyl, aryl, aralkyl, cycloalkyl, $-OR_{21}$, $-SO_2R_{21}$, $-COOR_{21}$, $-COR_{21}$, $-CONR_{21}R_{22}$, $R_{21}R_{22}P(O)$, $(R_{21}O)(R_{22}O)P(O)$, $R_{21}R_{22}P(S)$, $-SOR_{21}$ or $-Si(OR_{21})_3$, provided that when the two "R" groups on a carbon adjacent to the nitrogen are $=O$ or $=S$, A , B and C are hydrogen, alkyl, aralkyl or aryl and R_{21} and R_{22} are independently selected from hydrogen, alkyl, aryl, aralkyl or alkaryl groups.
- 25
- 30 2. The process of Claim 1 wherein said coupling agent is selected from the group consisting of phosphoryl azides, carbodiimides, cyanamides, ketenimines, isoxazolium salts, monocyclic nitrogen containing heterocyclic amides of aromatic character containing 1 to 4 nitrogens in the ring, alkoxylated acetylene, reagents which form a mixed anhydride with the carboxyl moiety of the dicarboxylic acid,
- 35 nitrogen containing heterocyclic compounds having a hydroxy group on one ring nitrogen or carbonates thereof in conjunction with a coupling agent from one of the

-60-

other groups, reagents which form an active ester with the carboxyl moiety of the dicarboxylic acid, benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate, re-dox based coupling agents, and mixtures thereof.

3. The process of Claim 2 wherein said coupling agent is
 5 diphenylphosphoryl azide or 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride and hydroxybenzotriazole.


4. The process of Claim 1 wherein said solvent is a polar organic solvent.

5. The process of Claim 1 wherein said base is selected from the group
 10 consisting of trialkylamines, pyridines and imidazoles.

6. The process of Claim 1 wherein R is hydrogen.

7. The process of Claim 1 wherein said diamine is represented by the formula:

15




8. The process of Claim 7 wherein R_{13} , R_{14} , R_{15} and R_{16} are
 20 independently selected from hydrogen or alkyl; or
 R_{13} or R_{14} and R_{15} or R_{16} together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms.

9. The process of Claim 8 wherein at least one of R_{13} , R_{14} , R_{15} and
 25 R_{16} is an alkyl group.

10. The process of Claim 9 wherein at least one of R_{13} and R_{14} and at least one of R_{15} and R_{16} is an alkyl group.

11. The process of Claim 8 wherein R_{13} and R_{16} together with the carbon atoms to which they are attached independently form a saturated, partially
 30 saturated or unsaturated cyclic having 3 to 20 carbon atoms and R_{14} and R_{15} are hydrogen.

12. The process of Claim 1 wherein said diamine is represented by the formula:




10

13. The process of Claim 12 wherein R_{13} , R_{14} , R_{15} , R_{16} , R_{19} and R_{20} are independently selected from hydrogen or alkyl; or R_{13} or R_{14} and R_{19} or R_{20} , or R_{15} or R_{16} and R_{19} or R_{20} together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms; or R_{19} and R_{20} together are $=\text{O}$ or $=\text{S}$.



14. The process of Claim 1 wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , R_9 , R_{10} , R_{11} and R_{12} are independently selected from hydrogen, alkyl radicals or radicals attached to the α -carbon of α -amino acids; or R_3 or R_4 and R_5 or R_6 , and R_7 or R_8 and R_9 or R_{10} together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms; or R_1 and R_2 , R_3 and R_4 , R_5 and R_6 , R_7 and R_8 , R_9 and R_{10} , and R_{11} and R_{12} independently are $=\text{O}$ or $=\text{S}$.

15. The process of Claim 14 wherein at least one of R_3 or R_4 and R_5 or R_6 , and R_7 or R_8 and R_9 or R_{10} together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms, and the remaining of R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , R_9 , R_{10} , R_{11} and R_{12} are independently selected from hydrogen, alkyl radicals or radicals attached to the α -carbon of α -amino acids.

16. The process of Claim 14 wherein at least one of R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , R_9 , R_{10} , R_{11} and R_{12} are alkyl radicals or radicals attached to the α -carbon of α -amino acids, and the remaining of R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , R_9 , R_{10} , R_{11} and R_{12} are hydrogen.



-63-

or
(Z)_n(Z)_n


and, optionally, further conducting an exchange reaction with said manganese complex to exchange the ligands X, Y and Z on said manganese complex;
 wherein X, Y and Z are ligands independently selected from the group consisting of
 halide, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxo, hydroperoxo,
 alkylperoxo, arylperoxo, ammonia, alkylamino, arylamino, heterocycloalkyl amino,
 heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine,
 nitric oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile,
 aryl nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid,
 aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic
 acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic
 acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol
 thiocarboxylic acid, alkyl carboxylic acid, aryl carboxylic acid, urea, alkyl urea, aryl
 urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea,

-64-

- sulfate, sulfite, bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate, thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl dithiocarbamate, aryl dithiocarbamate, alkylaryl dithiocarbamate, bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite, hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate, tetraaryl borate, tetraalkyl borate, tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino acid, hydroxamic acid, thiotosylate, and anions of ion exchange resins, or the corresponding anions thereof, or X, Y and Z are independently attached to one or more of the "R" groups and n is an integer from 0 to 3.


24. The process of Claim 23 wherein X, Y and Z are independently selected from the group consisting of halide, organic acid, nitrate and bicarbonate anions.

25. A process for preparing a substituted polyazamacrocyclic comprising contacting (a) a triamine represented by the formula:




- and (b) a dicarboxylic acid or ester thereof represented by the formula:

-65-



or an anhydride thereof; in the presence of a suitable base and a suitable solvent under
 5 reaction conditions sufficient to produce said substituted polyazamacrocyclic
 represented by the formula:



provided that when an ester of said dicarboxylic acid is used, said suitable base is
 20 optional and when said dicarboxylic acid or an anhydride of said dicarboxylic acid is
 used, the reaction mixture further comprises a suitable coupling agent;
 wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 and R_8 are independently selected from
 hydrogen or alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cycloalkylalkyl,
 cycloalkylcycloalkyl, cycloalkenylalkyl, alkylcycloalkyl, alkenylcycloalkyl,
 25 alkylcycloalkenyl, alkenylcycloalkenyl, heterocyclic, aryl and aralkyl radicals or
 radicals attached to the α -carbon of α -amino acids; or
 R_3 or R_4 and R_5 or R_6 together with the carbon atoms to which they are attached
 independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20
 carbon atoms; or

- R₁ or R₂ and R₃ or R₄, and R₅ or R₆ and R₇ or R₈ together with the carbon atoms to which they are attached independently form a nitrogen containing heterocycle having 2 to 20 carbon atoms provided that when the nitrogen containing heterocycle is an aromatic heterocycle which does not contain a substituent attached to the nitrogen, the hydrogen attached to the nitrogen in said formula, which nitrogen is also in the macrocycle and the R groups attached to the same carbon atoms of the macrocycle are absent;
- 5 R₁ and R₂, R₃ and R₄, R₅ and R₆, and R₇ and R₈ together with the carbon atom to which they are attached independently form a saturated, partially saturated or unsaturated ring structure having 3 to 20 carbon atoms or R₁ and R₂, R₃ and R₄, R₅ and R₆, and R₇ and R₈ independently are =O or =S; and combinations thereof; and wherein R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₉ and R₂₀ are independently selected from hydrogen or alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, alkylcycloalkyl, alkenylcycloalkyl, alkylcycloalkenyl, alkenylcycloalkenyl, heterocyclic, aryl and aralkyl radicals; or
- 10 R₉ or R₁₀ and R₁₁ or R₁₂, and R₁₃ or R₁₄ and R₁₅ or R₁₆ together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms, provided that if R₉ or R₁₀ and R₁₁ or R₁₂ together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic and the diamine has three carbons between the nitrogen atoms, the saturated, partially saturated or unsaturated cyclic has 4 to 20 carbon atoms; or
- 20 R₉ or R₁₀ and R₁₉ or R₂₀, or R₁₁ or R₁₂ and R₁₉ or R₂₀ together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms; or
- 25 R₁₁ or R₁₂ and R₁₃ or R₁₄ together with the carbon atoms to which they are attached independently form a nitrogen containing heterocycle having 2 to 20 carbon atoms provided that when the nitrogen containing heterocycle is an aromatic heterocycle which does not contain a substituent attached to the nitrogen, the hydrogen attached to the nitrogen in the formula, which nitrogen is also in the macrocycle and the R groups attached to the same carbon atoms of the macrocycle are absent;
- 30 R₉ and R₁₀, R₁₁ and R₁₂, R₁₃ and R₁₄, R₁₅ and R₁₆, and R₁₉ and R₂₀ together with the carbon atom to which they are attached independently form a saturated, partially saturated or unsaturated ring structure having 3 to 20 carbon atoms; or
- 35

-67-

R_{11} and R_{12} , and R_{13} and R_{14} independently are =O or =S; R_{19} and R_{20} independently are =O, =S, =NR₂₃, =N-OH, =N-OR₂₃, =N-O-C(O)-R₂₃, or =CR₂₃R₂₄; and combinations thereof;

wherein R_{17} and R_{18} are independently selected from hydrogen and alkyl groups;

5 wherein R is hydrogen or an alkyl group; and

wherein A, B and C are independently selected from hydrogen, alkyl, aryl, aralkyl, cycloalkyl, -OR₂₁, -SO₂R₂₁, -COOR₂₁, -COR₂₁, -CONR₂₁R₂₂, R₂₁R₂₂P(O), (R₂₁O)(R₂₂O)P(O), R₂₁R₂₂P(S), -SOR₂₁ or -Si(OR₂₁)₃, provided that when the two "R" groups on a carbon adjacent to the nitrogen are =O or =S, A, B and C are

10 hydrogen, alkyl, aralkyl or aryl, and R_{21} and R_{22} are independently selected from hydrogen, alkyl, aryl, aralkyl or alkaryl groups.

26. The process of Claim 25 wherein said coupling agent is selected from the group consisting of phosphoryl azides, carbodiimides, cyanamides, ketenimines, isoxazolium salts, monocyclic nitrogen containing heterocyclic amides
15 of aromatic character containing 1 to 4 nitrogens in the ring, alkoxyated acetylene, reagents which form a mixed anhydride with the carboxyl moiety of the dicarboxylic acid, nitrogen containing heterocyclic compounds having a hydroxy group on one ring nitrogen or carbonate thereof in conjunction with a coupling agent from one of the other groups, reagents which form an active ester with the carboxyl moiety of the
20 dicarboxylic acid, benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate, re-dox based coupling agents, and mixtures thereof.


27. The process of Claim 26 wherein said coupling agent is diphenylphosphoryl azide or 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride and hydroxybenzotriazole.

25 28. The process of Claim 25 wherein said solvent is a polar organic solvent.

29. The process of Claim 25 wherein said base is selected from the group consisting of trialkylamines, pyridines and imidazoles.

30. The process of Claim 25 wherein R is hydrogen.

30 31. The process of Claim 25 wherein said triamine is represented by the formula:




32. The process of Claim 31 wherein R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅ and R₁₆ are independently selected from hydrogen or alkyl; or R₉ or R₁₀ and R₁₁ or R₁₂, and R₁₃ or R₁₄ and R₁₅ or R₁₆ together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms; or R₁₁ and R₁₂, and R₁₃ and R₁₄ are independently =O or =S.

33. The process of Claim 32 wherein at least one of R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅ and R₁₆ is an alkyl group.

34. The process of Claim 33 wherein at least one of R₁₃ and R₁₄ and at least one of R₁₅ and R₁₆ is an alkyl group or at least one of R₉ and R₁₀ and at least one of R₁₁ and R₁₂ is an alkyl group.

35. The process of Claim 32 wherein R₉ and R₁₂ or R₁₃ and R₁₆ together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms and the remaining R groups of said triamine are hydrogen.

36. The process of Claim 25 wherein said triamine is represented by the formula:



25

37. The process of Claim 36 wherein R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₉ and R₂₀ are independently selected from hydrogen or alkyl; or R₉ or R₁₀ and R₁₉ or R₂₀, R₁₁ or R₁₂ and R₁₉ or R₂₀, or R₁₃ or R₁₄ and R₁₅ or R₁₆ together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms; or R₁₁ and R₁₂, R₁₃ and R₁₄, and R₁₉ and R₂₀ independently are =O or =S.

38. The process of Claim 25 wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇ and R₈ are independently selected from hydrogen, alkyl radicals or radicals attached to the α -carbon of α -amino acids; or

-69-

R_3 or R_4 and R_5 or R_6 together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms; or

R_1 and R_2 , R_3 and R_4 , R_5 and R_6 , and R_7 and R_8 independently are $=O$ or $=S$.

- 5 39. The process of Claim 38 wherein R_3 or R_4 and R_5 or R_6 together with the carbon atoms to which they are attached independently form a saturated, partially saturated or unsaturated cyclic having 3 to 20 carbon atoms, and the remaining of R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 and R_8 are independently selected from hydrogen, alkyl radicals or radicals attached to the α -carbon of α -amino acids.


- 10 40. The process of Claim 38 wherein at least one of R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 and R_8 are alkyl radicals or radicals attached to the α -carbon of α -amino acids, and the remaining of R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 and R_8 are hydrogen.

41. The process of Claim 38 wherein R is hydrogen.

- 15 42. The process of Claim 38 wherein A , B and C are independently selected from the group consisting of hydrogen, alkyl, aryl and aralkyl.

43. The process of Claim 25 wherein R_{17} and R_{18} are hydrogen.

44. The process of Claim 25 further comprising reducing said substituted polyazamacrocyclic to produce a reduced substituted polyazamacrocyclic of the formula:




wherein A , B and C are independently selected from hydrogen, alkyl, aryl, aralkyl and cycloalkyl groups.


-70-

45. The process of Claim 44 wherein said reducing agent is selected
 5 from the group consisting of lithium aluminum hydride, borane and sodium bis(2-methoxyethoxy) aluminum hydride.

46. The process of Claim 43 further comprising reacting said reduced
 substituted polyazamacrocyclic with a manganese compound under essentially
 anhydrous and anaerobic conditions to produce a manganese complex represented by
 10 the formula:



or



SUBSTITUTE SHEET
 ISA/EP

-71-

- and, optionally, further conducting an exchange reaction with said manganese complex to exchange the ligands X, Y and Z on said manganese complex;
 wherein X, Y and Z are ligands independently selected from the group consisting of halide, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxy, hydroperoxy,
 5 alkylperoxy, arylperoxy, ammonia, alkylamino, arylamino, heterocycloalkyl amino, heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine, nitric oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol thiocarboxylic acid, alkyl carboxylic acid, aryl carboxylic acid, urea, alkyl urea, aryl urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea, sulfate, sulfite, bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl
 10 phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate, thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen
 15 phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl dithiocarbamate, aryl dithiocarbamate, alkylaryl dithiocarbamate, bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite, hypobromite,
 20 tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate, tetraaryl borate, tetraalkyl borate, tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino acid, hydroxamic acid, thiotosylate, and anions of ion exchange resins, or the corresponding anions thereof, or X, Y and Z are independently attached to one or more of the "R" groups and n is an integer from 0 to
 30 3.

47. The process of Claim 46 wherein X, Y and Z are independently selected from the group consisting of halide, organic acid, nitrate and bicarbonate anions.

48. The process of Claim 44 wherein said reduction is conducted in the presence of a reducing agent selected from the group consisting of aluminum hydrides and boron hydrides.

SUBSTITUTE SHEET
ISA/EP

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 96/07553

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D259/00 C07F9/6524 C07F13/00 C07F15/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D C07F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO,A,92 08707 (SALUTAR) 29 May 1992 see page 16 - page 17 see page 31; example 1 see page 44 - page 54; claims ---	1-47
A	WO,A,94 15925 (MONSANTO COMPANY) 21 July 1994 see page 54 - page 59 see page 23 ---	1-47
A	EP,A,0 524 161 (MONSANTO COMPANY) 20 January 1993 see page 71 - page 78; claims ---	20-23, 44-47
P,A	WO,A,95 28968 (MONSANTO COMPANY) 2 November 1995 see the whole document -----	1-47

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

3 September 1996

Date of mailing of the international search report

10.09.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Luyten, H

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/US 96/07553

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9208707	29-05-92	AT-T- 140219 AU-B- 8862891 DE-D- 69120826 EP-A- 0556240 JP-T- 6504301	15-07-96 11-06-92 14-08-96 25-08-93 19-05-94
WO-A-9415925	21-07-94	AU-B- 5964894 CA-A- 2151262 EP-A- 0679155 JP-T- 8505622	15-08-94 21-07-94 02-11-95 18-06-96
EP-A-524161	20-01-93	AU-B- 661023 AU-B- 2338392 CA-A- 2072897 CA-A- 2072934 EP-A- 0598753 JP-T- 6509566 NZ-A- 272364 WO-A- 9302090 ZA-A- 9205139	13-07-95 23-02-93 20-01-93 20-01-93 01-06-94 27-10-94 27-02-96 04-02-93 26-04-93
WO-A-9528968	02-11-95	AU-B- 2128895	16-11-95